

Some remarks about the second Leibniz cohomology group for Lie algebras

L. Magnin

Institut Mathématique de Bourgogne *
magnin@u-bourgogne.fr

December 15, 2008

Abstract

We compare by a very elementary approach the second adjoint and trivial Leibniz cohomology spaces of a Lie algebra to the usual ones. Examples are given of coupled cocycles. Some properties are deduced as to Leibniz deformations. We also consider the class of Lie algebras for which the Koszul 3-form is zero, and prove that it contains all quotients of Borel subalgebras, or of their nilradicals, of finite dimensional semisimple Lie algebras. Finally, a list of Kac-Moody types for indecomposable nilpotent Lie algebras of dimension ≤ 7 is given.

1 Introduction.

Leibniz algebras, along with their Leibniz cohomologies, were introduced in [11] as a non antisymmetric version of Lie algebras. Lie algebras are special Leibniz algebras, and Pirashvili introduced [19] a spectral sequence, that, when applied to Lie algebras, measures the difference between the Lie algebra cohomology and the Leibniz cohomology. Now, Lie algebras have deformations as Leibniz algebras and those are piloted by the adjoint Leibniz 2-cocycles. In the present paper, we focus on the second Leibniz cohomology groups $HL^2(\mathfrak{g}, \mathfrak{g})$, $HL^2(\mathfrak{g}, \mathbb{C})$ for adjoint and trivial representations of a complex Lie algebra $\mathfrak{g}$. We adopt a very elementary approach, not resorting to the Pirashvili sequence, to compare $HL^2(\mathfrak{g}, \mathfrak{g})$ and $HL^2(\mathfrak{g}, \mathbb{C})$ to $H^2(\mathfrak{g}, \mathfrak{g})$ and $H^2(\mathfrak{g}, \mathbb{C})$ respectively. In both cases, HL^2 appears to be the direct sum of 3 spaces: $H^2 \oplus ZL_0^2 \oplus \mathcal{C}$ where H^2 is the Lie algebra cohomology group, ZL_0^2 is the space of symmetric Leibniz-2-cocycles and $\mathcal{C}$ is a space of *coupled* Leibniz-2-cocycles the nonzero elements of which have the property that their symmetric and antisymmetric parts are not Leibniz cocycles (a similar situation occurs in second cohomology of current algebras [23]). Many Lie algebras are uncoupling in the sense that the space of

*UMR CNRS 5584, Université de Bourgogne, BP 47870, 21078 Dijon Cedex, France.

coupled cocycles is zero. We give examples in dimensions starting from 5 of Lie algebras that are not uncoupling. We deduce some criteria for a Lie algebra to be rigid as Leibniz algebra, and results as to the number of Leibniz deformations in some cases. In a first part, we consider the subclass of uncoupling Lie algebras which have the property that the Koszul 3-form is zero. We prove in particular that it contains all quotients of Borel subalgebras, or of their nilradicals, of finite dimensional semisimple Lie algebras. We also give a list of Kac-Moody types for indecomposable nilpotent Lie algebras of dimension ≤ 7 .

2 Leibniz cohomology.

Recall that a (right) Leibniz algebra is an algebra $\mathfrak{g}$ with a (non necessarily anti-symmetric) bracket, such that the right adjoint operations $[\cdot, Z]$ are required to be derivations for any $Z \in \mathfrak{g}$. In the presence of antisymmetry, that is equivalent to the Jacobi identity, hence any Lie algebra is a Leibniz algebra.

The Leibniz cohomology $HL^\bullet(\mathfrak{g}, \mathfrak{g})$ of a Leibniz algebra is defined from the complex $CL^\bullet(\mathfrak{g}, \mathfrak{g}) = \text{Hom}(\mathfrak{g}^{\otimes \bullet}, \mathfrak{g}) = \mathfrak{g} \otimes (\mathfrak{g}^*)^{\otimes \bullet}$ with the Leibniz-coboundary δ defined for $\psi \in CL^n(\mathfrak{g}, \mathfrak{g})$ by

$$\begin{aligned} (\delta\psi)(X_1, X_2, \dots, X_{n+1}) = \\ [X_1, \psi(X_2, \dots, X_{n+1})] + \sum_{i=2}^{n+1} (-1)^i [\psi(X_1, \dots, \hat{X}_i, \dots, X_{n+1}), X_i] \\ + \sum_{1 \leq i < j \leq n+1} (-1)^{j+1} \psi(X_1, \dots, X_{i-1}, [X_i, X_j], X_{i+1}, \dots, \hat{X}_j, \dots, X_{n+1}). \end{aligned}$$

(If $\mathfrak{g}$ is a Lie algebra, δ coincides with the usual coboundary d on $C^\bullet(\mathfrak{g}, \mathfrak{g}) = \mathfrak{g} \otimes \bigwedge^\bullet \mathfrak{g}^*$.)

For $\psi \in CL^1(\mathfrak{g}, \mathfrak{g}) = C^1(\mathfrak{g}, \mathfrak{g}) = \mathfrak{g} \otimes \mathfrak{g}^*$

$$(\delta\psi)(X, Y) = [X, \psi(Y)] + [\psi(X), Y] - \psi([X, Y]).$$

For $\psi \in CL^2(\mathfrak{g}, \mathfrak{g}) = \mathfrak{g} \otimes (\mathfrak{g}^*)^{\otimes 2}$,

$$\begin{aligned} (\delta\psi)(X, Y, Z) = [X, \psi(Y, Z)] + [\psi(X, Z), Y] - [\psi(X, Y), Z] \\ - \psi([X, Y], Z) + \psi(X, [Y, Z]) + \psi([X, Z], Y). \end{aligned}$$

In the same way, the trivial Leibniz cohomology $HL^\bullet(\mathfrak{g}, \mathbb{C})$ is defined from the complex $CL^\bullet(\mathfrak{g}, \mathbb{C}) = (\mathfrak{g}^*)^{\otimes \bullet}$ with the trivial-Leibniz-coboundary $\delta_{\mathbb{C}}$ defined for $\psi \in CL^n(\mathfrak{g}, \mathbb{C})$ by

$$\begin{aligned} (\delta_{\mathbb{C}}\psi)(X_1, X_2, \dots, X_{n+1}) = \\ \sum_{1 \leq i < j \leq n+1} (-1)^{j+1} \psi(X_1, \dots, X_{i-1}, [X_i, X_j], X_{i+1}, \dots, \hat{X}_j, \dots, X_{n+1}). \end{aligned}$$

If $\mathfrak{g}$ is a Lie algebra, $\delta_{\mathbb{C}}$ is the usual coboundary $d_{\mathbb{C}}$ on $C^{\bullet}(\mathfrak{g}, \mathbb{C}) = \bigwedge^{\bullet} \mathfrak{g}^*$.
For $\psi \in CL^1(\mathfrak{g}, \mathbb{C}) = \mathfrak{g}^* \otimes \mathfrak{g}$,

$$(\delta_{\mathbb{C}}\psi)(X, Y) = -\psi([X, Y]).$$

For $\psi \in CL^2(\mathfrak{g}, \mathbb{C}) = (\mathfrak{g}^*)^{\otimes 2}$,

$$(\delta_{\mathbb{C}}\psi)(X, Y, Z) = -\psi([X, Y], Z) + \psi(X, [Y, Z]) + \psi([X, Z], Y).$$

3 Some properties of the Koszul map.

Let $\mathfrak{g}$ be any finite dimensional complex Lie algebra. Recall that a symmetric bilinear form $B \in S^2 \mathfrak{g}^*$ is invariant, i.e. $B \in (S^2 \mathfrak{g}^*)^{\mathfrak{g}}$ if and only if $B([Z, X], Y) = -B(X, [Z, Y]) \forall X, Y, Z \in \mathfrak{g}$. The Koszul map [9] $\mathcal{I} : (S^2 \mathfrak{g}^*)^{\mathfrak{g}} \rightarrow (\bigwedge^3 \mathfrak{g}^*)^{\mathfrak{g}} \subset Z^3(\mathfrak{g}, \mathbb{C})$ is defined by $\mathcal{I}(B) = I_B$, with $I_B(X, Y, Z) = B([X, Y], Z) \forall X, Y, Z \in \mathfrak{g}$.

Lemma 1. Denote $\mathcal{C}^2 \mathfrak{g} = [\mathfrak{g}, \mathfrak{g}]$. The projection $\pi : \mathfrak{g} \rightarrow \mathfrak{g}/\mathcal{C}^2 \mathfrak{g}$ induces an isomorphism

$$\varpi : \ker \mathcal{I} \rightarrow S^2 (\mathfrak{g}/\mathcal{C}^2 \mathfrak{g})^*.$$

Proof. For $B \in \ker \mathcal{I}$, define $\varpi(B) \in S^2 (\mathfrak{g}/\mathcal{C}^2 \mathfrak{g})^*$ by

$$\varpi(B)(\pi(X), \pi(Y)) = B(X, Y), \forall X, Y \in \mathfrak{g}.$$

$\varpi(B)$ is well-defined since for $X, Y, U, V \in \mathfrak{g}$

$$\begin{aligned} B(X + [U, V], Y) &= B(X, Y) + B([U, V], Y) \\ &= B(X, Y) + I_B(U, V, Y) \\ &= B(X, Y) \text{ (as } I_B = 0). \end{aligned}$$

The map ϖ is injective since $\varpi(B) = 0$ implies $B(X, Y) = 0 \forall X, Y \in \mathfrak{g}$. To prove that it is onto, let $\bar{B} \in S^2 (\mathfrak{g}/\mathcal{C}^2 \mathfrak{g})^*$, and let $B_{\pi} \in S^2 \mathfrak{g}^*$ defined by $B_{\pi}(X, Y) = \bar{B}(\pi(X), \pi(Y))$. Then $B_{\pi}([X, Y], Z) = \bar{B}(\pi([X, Y]), \pi(Z)) = \bar{B}(0, \pi(Z)) = 0 \forall X, Y, Z \in \mathfrak{g}$, hence $B_{\pi} \in (S^2 \mathfrak{g}^*)^{\mathfrak{g}}$ and $B_{\pi} \in \ker \mathcal{I}$. Now, $\varpi(B_{\pi}) = \bar{B}$. $\square$

From lemma 1, $\dim (S^2 \mathfrak{g}^*)^{\mathfrak{g}} = \frac{p(p+1)}{2} + \dim \text{Im } \mathcal{I}$, where $p = \dim H^1(\mathfrak{g}, \mathbb{C})$. For reductive $\mathfrak{g}$, $\dim (S^2 \mathfrak{g}^*)^{\mathfrak{g}} = \dim H^3(\mathfrak{g}, \mathbb{C})$ ([9]). Note also that the restriction of $\delta_{\mathbb{C}}$ to $(S^2 \mathfrak{g}^*)^{\mathfrak{g}}$ is $-\mathcal{I}$.

Definition 1. $\mathfrak{g}$ is said to be $\mathcal{I}$ -null (resp. $\mathcal{I}$ -exact) if $\mathcal{I} = 0$ (resp. $\text{Im } \mathcal{I} \subset B^3(\mathfrak{g}, \mathbb{C})$).

$\mathfrak{g}$ is $\mathcal{I}$ -null if and only $\mathcal{C}^2 \mathfrak{g} \subset \ker B \forall B \in (S^2 \mathfrak{g}^*)^{\mathfrak{g}}$. It is standard that for any $B \in (S^2 \mathfrak{g}^*)^{\mathfrak{g}}$, there exists $B_1 \in (S^2 \mathfrak{g}^*)^{\mathfrak{g}}$ such that $\ker (B + B_1) \subset \mathcal{C}^2 \mathfrak{g}$, hence $\bigcap_{B \in (S^2 \mathfrak{g}^*)^{\mathfrak{g}}} \ker B \subset \mathcal{C}^2 \mathfrak{g}$, hence $\mathfrak{g}$ is $\mathcal{I}$ -null if and only $\bigcap_{B \in (S^2 \mathfrak{g}^*)^{\mathfrak{g}}} \ker B = \mathcal{C}^2 \mathfrak{g}$.

Lemma 2. (i) Any quotient of a (not necessarily finite dimensional) $\mathcal{I}$ -null Lie algebra is $\mathcal{I}$ -null.

(ii) any finite direct product of $\mathcal{I}$ -null Lie algebras is $\mathcal{I}$ -null.

Proof. (i) Let $\mathfrak{g}$ be any $\mathcal{I}$ -null Lie algebra, $\mathfrak{h}$ an ideal of $\mathfrak{g}$, $\bar{\mathfrak{g}} = \mathfrak{g}/\mathfrak{h}$, $\pi : \mathfrak{g} \rightarrow \bar{\mathfrak{g}}$ the projection, and $\bar{B} \in (S^2 \bar{\mathfrak{g}}^*)^{\bar{\mathfrak{g}}}$. Define $B_\pi \in S^2 \mathfrak{g}^*$ by $B_\pi(X, Y) = \bar{B}(\pi(X), \pi(Y))$, $X, Y \in \mathfrak{g}$. Then $B_\pi([X, Y], Z) = \bar{B}(\pi([X, Y]), \pi(Z)) = \bar{B}([\pi(X), \pi(Y)], \pi(Z)) = \bar{B}(\pi(X), [\pi(Y), \pi(Z)]) = \bar{B}(\pi(X), \pi([Y, Z])) = B_\pi(X, [Y, Z]) \forall X, Y, Z \in \mathfrak{g}$, hence $B_\pi \in (S^2 \mathfrak{g}^*)^{\mathfrak{g}}$ and $I_{B_\pi} \circ (\pi \times \pi \times \pi) = I_{\bar{B}} = 0$ since $\bar{\mathfrak{g}}$ is $\mathcal{I}$ -null. Hence $I_{B_\pi} = 0$.

(ii) Let $\mathfrak{g} = \mathfrak{g}_1 \times \mathfrak{g}_2$ ($\mathfrak{g}_1, \mathfrak{g}_2$ $\mathcal{I}$ -null) and $B \in (S^2 \mathfrak{g}^*)^{\mathfrak{g}}$. As $B(X_1, [Y_2, Z_2]) = B([X_1, Y_2], Z_2) = B(0, Z_2) = 0 \forall X_1 \in \mathfrak{g}_1, Y_2, Z_2 \in \mathfrak{g}_2$, B vanishes on $\mathfrak{g}_1 \times \mathcal{C}^2 \mathfrak{g}_2$ and on $\mathcal{C}^2 \mathfrak{g}_1 \times \mathfrak{g}_2$ as well, hence $I_B = 0$. $\square$

Lemma 3. Let $\mathfrak{g}$ be a finite dimensional semi-simple Lie algebra, with Cartan subalgebra $\mathfrak{h}$, simple root system S , positive roots Δ_+ , and root subspaces $\mathfrak{g}^\alpha$. Let $\mathfrak{k}$ be any non empty subspace of $\mathfrak{h}$, and $\Gamma \subset \Delta_+$ such that $\alpha + \beta \in \Gamma$ for $\alpha, \beta \in \Gamma$, $\alpha + \beta \in \Delta_+$. Consider $\mathfrak{u} = \mathfrak{k} \oplus \bigoplus_{\alpha \in \Gamma} \mathfrak{g}^\alpha$.

(i) Suppose that $\alpha|_{\mathfrak{k}} \neq 0 \forall \alpha \in \Gamma$. Then $\mathfrak{u}$ is $\mathcal{I}$ -null.

(ii) Suppose that $\alpha|_{\mathfrak{k}} = 0 \forall \alpha \in \Gamma \cap S$, and $\alpha|_{\mathfrak{k}} \neq 0 \forall \alpha \in \Gamma \setminus S$. Then $\mathfrak{u}$ is $\mathcal{I}$ -null.

Proof. (i) Let $\mathfrak{u}_+ = \bigoplus_{\alpha \in \Gamma} \mathfrak{g}^\alpha$, and X_α a root vector in $\mathfrak{g}^\alpha$: $\mathfrak{g}^\alpha = \mathbb{C}X_\alpha \forall \alpha \in \Gamma$. Let $B \in (S^2 \mathfrak{u}^*)^{\mathfrak{u}}$. First, $B(H, X) = 0 \forall H \in \mathfrak{k}, X \in \mathfrak{u}_+$. In fact, for any $\alpha \in \Gamma$, since there exists $H_\alpha \in \mathfrak{k}$ such that $\alpha(H_\alpha) \neq 0$, $B(H, X_\alpha) = \frac{1}{\alpha(H_\alpha)} B(H, [H_\alpha, X_\alpha]) = \frac{1}{\alpha(H_\alpha)} B([H, H_\alpha], X_\alpha) = \frac{1}{\alpha(H_\alpha)} B(0, X_\alpha) = 0$. Second, that entails that the restriction of B to $\mathfrak{u}_+ \times \mathfrak{u}_+$ is zero, since for any $\alpha, \beta \in \Gamma$,

$$B(X_\alpha, X_\beta) = \frac{1}{\alpha(H_\alpha)} B([H_\alpha, X_\alpha], X_\beta) = \frac{1}{\alpha(H_\alpha)} B(H_\alpha, [X_\alpha, X_\beta]) = 0$$

as $[X_\alpha, X_\beta] \in \mathfrak{u}_+$. Then $\mathfrak{u}$ is $\mathcal{I}$ -null.

(ii) In that case, $X_\alpha \notin \mathcal{C}^2 \mathfrak{u} \forall \alpha \in \Gamma \cap S$, and $\dim(\mathfrak{u} / \mathcal{C}^2 \mathfrak{u}) = \dim \mathfrak{k} + \#(\Gamma \cap S)$. For $\mathfrak{u}$ to be $\mathcal{I}$ -null, one has to prove that, for any $B \in (S^2 \mathfrak{u}^*)^{\mathfrak{u}}$:

$$B(H, X_\beta) = 0 \forall H \in \mathfrak{k}, \beta \in \Gamma \setminus S; \quad (1)$$

$$B(X_\alpha, X_\beta) = 0 \forall \alpha \in \Gamma \cap S, \beta \in \Gamma \setminus S; \quad (2)$$

$$B(X_\beta, X_\gamma) = 0 \forall \alpha, \beta \in \Gamma \setminus S. \quad (3)$$

(1) is proved as in case (i). To prove (2), let $H_\beta \in \mathfrak{k}$ such that $\beta(H_\beta) \neq 0$. Then

$$\begin{aligned} B(X_\alpha, X_\beta) &= \frac{1}{\beta(H_\beta)} B(X_\alpha, [H_\beta, X_\beta]) = \frac{1}{\beta(H_\beta)} B([X_\alpha, H_\beta], X_\beta) = \\ &= -\frac{1}{\beta(H_\beta)} B(\alpha(H_\beta)X_\alpha, X_\beta) = -\frac{1}{\beta(H_\beta)} B(0, X_\beta) = 0. \end{aligned}$$

As to (3),

$$B(X_\beta, X_\gamma) = \frac{1}{\beta(H_\beta)} B([H_\beta, X_\beta], X_\gamma) = \frac{1}{\beta(H_\beta)} B(H_\beta, [X_\beta, X_\gamma]) = 0 \text{ from (1).}$$

□

Example 1. Any Borel subalgebra is $\mathcal{I}$ -null.

Proposition 1. Let $\mathfrak{g}_2$ be a codimension 1 ideal of the Lie algebra $\mathfrak{g}$, $(x_1, \dots, x_N)$ a basis of $\mathfrak{g}$ with $x_1 \notin \mathfrak{g}_2$, $x_2, \dots, x_N \in \mathfrak{g}_2$, π_2 the corresponding projection onto $\mathfrak{g}_2$, and $(\omega^1, \dots, \omega^N)$ denote the dual basis for $\mathfrak{g}^*$. Let $B \in (S^2 \mathfrak{g}^*)^{\mathfrak{g}}$, and denote $B_2 \in (S^2 \mathfrak{g}_2^*)^{\mathfrak{g}_2}$ the restriction of B to $\mathfrak{g}_2 \times \mathfrak{g}_2$. Then:

(i)

$$I_B = d(\omega^1 \wedge f) + I_{B_2} \circ (\pi_2 \times \pi_2 \times \pi_2). \quad (4)$$

where $f = B(\cdot, x_1) \in \mathfrak{g}^*$.

(ii) Let $\gamma \in \bigwedge^2 \mathfrak{g}_2^* \subset \bigwedge^2 \mathfrak{g}^*$, and denote $d_{\mathfrak{g}_2}$ the coboundary operator of $\mathfrak{g}_2$. Then

$$d\gamma = \omega^1 \wedge \theta_{x_1}(\gamma) + d_{\mathfrak{g}_2} \gamma \circ (\pi_2 \times \pi_2 \times \pi_2) \quad (5)$$

where θ_{x_1} stands for the coadjoint action of x_1 on the cohomology of $\mathfrak{g}$.

(iii) Suppose $I_{B_2} \in B^3(\mathfrak{g}_2, \mathbb{C})$, and let $\gamma \in \bigwedge^2 \mathfrak{g}_2^* \subset \bigwedge^2 \mathfrak{g}^*$ such that $I_{B_2} = d_{\mathfrak{g}_2} \gamma$. Then $I_B \in B^3(\mathfrak{g}, \mathbb{C})$ if and only if $\omega^1 \wedge \theta_{x_1}(\gamma) \in B^3(\mathfrak{g}, \mathbb{C})$. In particular, the condition

$$\theta_{x_1}(\gamma) = df \quad (6)$$

implies $I_B = d\gamma$.

Proof. (i) For $X, Y, Z \in \mathfrak{g}$ one has

$$\begin{aligned} B([X, Y], Z) &= B([\omega^1(X)x_1 + \pi_2(X), \omega^1(Y)x_1 + \pi_2(Y)], \omega^1(Z)x_1 + \pi_2(Z)) \\ &= B(\omega^1(X)[x_1, \pi_2(Y)] - \omega^1(Y)[x_1, \pi_2(X)] + [\pi_2(X), \pi_2(Y)], \\ &\quad \omega^1(Z)x_1 + \pi_2(Z)) \\ &= \omega^1(X)\omega^1(Z)B([x_1, \pi_2(Y)], x_1) - \omega^1(Y)\omega^1(Z)B([x_1, \pi_2(X)], x_1) \\ &\quad + \beta(X, Y, Z) + B([\pi_2(X), \pi_2(Y)], \pi_2(Z)) \\ &= \beta(X, Y, Z) + B([\pi_2(X), \pi_2(Y)], \pi_2(Z)) \end{aligned} \quad (7)$$

where

$$\begin{aligned} \beta(X, Y, Z) &= \omega^1(Z)B([\pi_2(X), \pi_2(Y)], x_1) + \omega^1(X)B([x_1, \pi_2(Y)], \pi_2(Z)) \\ &\quad - \omega^1(Y)B([x_1, \pi_2(X)], \pi_2(Z)) \\ &= \omega^1(Z)B([\pi_2(X), \pi_2(Y)], x_1) + \omega^1(X)B(x_1, [\pi_2(Y), \pi_2(Z)]) \\ &\quad - \omega^1(Y)B(x_1, [\pi_2(X), \pi_2(Z)]). \end{aligned}$$

Now

$$\begin{aligned} df(X, Y) &= -B([X, Y], x_1) \\ &= -B([\omega^1(X)x_1 + \pi_2(X), \omega^1(Y)x_1 + \pi_2(Y)], x_1) \\ &= -B(\omega^1(X)[x_1, \pi_2(Y)] - \omega^1(Y)[x_1, \pi_2(X)] + [\pi_2(X), \pi_2(Y)], x_1) \\ &= -B([\pi_2(X), \pi_2(Y)], x_1), \end{aligned}$$

hence

$$\begin{aligned}\beta(X, Y, Z) &= -(\omega^1(Z)df(X, Y) + \omega^1(X)df(Y, Z) - \omega^1(Y)df(X, Z)) \\ &= -(\omega^1 \wedge df)(X, Y, Z).\end{aligned}$$

Since $d\omega^1 = 0$, (7) then reads

$$I_B = d(\omega^1 \wedge f) + I_{B_2} \circ (\pi_2 \times \pi_2 \times \pi_2). \quad (8)$$

(ii) One has for any $X, Y, Z \in \mathfrak{g}$

$$\begin{aligned}d\gamma(X, Y, Z) &= d\gamma(\pi_2(X), \pi_2(Y), \pi_2(Z)) + \omega^1(X)d\gamma(x_1, \pi_2(Y), \pi_2(Z)) \\ &\quad + \omega^1(Y)d\gamma(\pi_2(X), x_1, \pi_2(Z)) + \omega^1(Z)d\gamma(\pi_2(X), \pi_2(Y), x_1).\end{aligned}$$

Now, since γ vanishes if one of its arguments is x_1 ,

$$\begin{aligned}d\gamma(x_1, \pi_2(Y), \pi_2(Z)) &= -\gamma([x_1, \pi_2(Y)], \pi_2(Z)) + \gamma([x_1, \pi_2(Z)], \pi_2(Y)) \\ d\gamma(\pi_2(X), x_1, \pi_2(Z)) &= -\gamma([\pi_2(X), x_1], \pi_2(Z)) - \gamma([x_1, \pi_2(Z)], \pi_2(X)) \\ d\gamma(\pi_2(X), \pi_2(Y), x_1) &= \gamma([\pi_2(X), x_1], \pi_2(Y)) - \gamma([\pi_2(Y), x_1], \pi_2(X)),\end{aligned}$$

hence

$$\begin{aligned}d\gamma(X, Y, Z) &= d\gamma(\pi_2(X), \pi_2(Y), \pi_2(Z)) + \omega^1(X)\theta_{x_1}\gamma(\pi_2(Y), \pi_2(Z)) \\ &\quad - \omega^1(Y)\theta_{x_1}\gamma(\pi_2(X), \pi_2(Z)) + \omega^1(Z)\theta_{x_1}\gamma(\pi_2(X), \pi_2(Y)) \\ &= d\gamma(\pi_2(X), \pi_2(Y), \pi_2(Z)) + (\omega^1 \wedge \theta_{x_1}\gamma)(X, Y, Z)\end{aligned}$$

since $\theta_{x_1}\gamma(\pi_2(U), \pi_2(V)) = \theta_{x_1}\gamma(U, V)$ for all $U, V \in \mathfrak{g}$.

(iii) Results immediately from (i) and (ii). $\square$

Corollary 1. *Under the hypotheses of Prop. 1, suppose that x_1 commutes with every x_i ($2 \leq i \leq N$) except for $x_{i_1}, \dots, x_{i_r}$ and that $x_{i_1}, \dots, x_{i_r}$ commute to one another. Then, if $\mathfrak{g}_2$ is $\mathcal{I}$ -null, $\mathfrak{g}$ is $\mathcal{I}$ -null.*

Proof. One has to prove that for any invariant bilinear symmetric form B on $\mathfrak{g}$, and any $2 \leq i, j \leq N$, $B(x_1, [x_i, x_j]) = 0$. For $i \neq i_1, \dots, i_r$, and any $j \geq 2$, $B(x_1, [x_i, x_j]) = B([x_1, x_i], x_j) = B(0, x_j) = 0$. For $i, j \in \{i_1, \dots, i_r\}$, $B(x_1, [x_i, x_j]) = B(x_1, 0) = 0$. $\square$

Corollary 2. *Any nilpotent standard filiform Lie algebra or any Heisenberg Lie algebra is $\mathcal{I}$ -null.*

Corollary 3. *Any Lie algebra containing some $\mathcal{I}$ -null codimension 1 ideal is $\mathcal{I}$ -exact.*

Corollary 4. *Any nilradical $\mathfrak{g}$ of a Borel subalgebra of a finite-dimensional semi-simple Lie algebra is $\mathcal{I}$ -null.*

Proof. It is enough to consider the case of a simple Lie algebra, hence of one of the 4 classical types plus the 5 exceptional ones.

Case A_n . Denote $E_{i,j}, 1 \leq i, j \leq n+1$ the canonical basis of $\mathfrak{gl}(n+1, \mathbb{C})$. One may suppose that the Borel subalgebra of A_n is comprised of the upper triangular matrices with zero trace, and the Cartan subalgebra $\mathfrak{h}$ is $\bigoplus_{i=1}^{i=n} \mathbb{C}H_i$ with $H_i = E_{i,i} - E_{i+1,i+1}$. For $n=1$, $\mathfrak{g}$ is $\mathbb{C}$, hence $\mathcal{I}$ -null. Suppose the result holds true for the nilradical of the Borel subalgebra of $A_{n-1} = \mathfrak{sl}(n)$. One has $\mathfrak{g} = \mathbb{C}E_{1,2} \oplus \cdots \oplus \mathbb{C}E_{1,n+1} \oplus \mathfrak{g}'_2$ with $\mathfrak{g}'_2$ having basis $E_{i,j}, 2 \leq i < j \leq n+1$ and being the nilradical of the Borel subalgebra of A_{n-1} , hence $\mathcal{I}$ -null. $E_{1,n+1}$ commutes with $\mathfrak{g}'_2$, hence, from corr. 1, $\mathbb{C}E_{1,n+1} \oplus \mathfrak{g}'_2$ is $\mathcal{I}$ -null. Now $E_{1,n}$ commutes with all members of the basis of $\mathbb{C}E_{1,n+1} \oplus \mathfrak{g}'_2$, except for $E_{n,n+1}$. Hence, from corr. 1, $\mathbb{C}E_{1,n} \oplus \mathbb{C}E_{1,n+1} \oplus \mathfrak{g}'_2$ is $\mathcal{I}$ -null. Now consider $\mathbb{C}E_{1,n-1} \oplus \mathbb{C}E_{1,n} \oplus \mathbb{C}E_{1,n+1} \oplus \mathfrak{g}'_2$. $\mathbb{C}E_{1,n-1}$ commutes with all members of the basis of $\mathbb{C}E_{1,n} \oplus \mathbb{C}E_{1,n+1} \oplus \mathfrak{g}'_2$ except for $E_{n-1,n}, E_{n-1,n+1}$. Since those 2 commute, we get in the same way that $\mathbb{C}E_{1,n-1} \oplus \mathbb{C}E_{1,n} \oplus \mathbb{C}E_{1,n+1} \oplus \mathfrak{g}'_2$ is $\mathcal{I}$ -null. The result then follows by induction.

Case D_n . We may take D_n as the Lie algebra of matrices

$$\begin{pmatrix} Z_1 & Z_2 \\ Z_3 & -{}^t Z_1 \end{pmatrix} \quad (9)$$

with $Z_i \in \mathfrak{gl}(n, \mathbb{C})$, Z_2, Z_3 skew symmetric (see [7], p. 193). Denote $\tilde{E}_{i,j} = \begin{pmatrix} E_{i,j} & 0 \\ 0 & -E_{j,i} \end{pmatrix}$, $\tilde{F}_{i,j} = \begin{pmatrix} 0 & E_{i,j} - E_{j,i} \\ 0 & 0 \end{pmatrix}$ ($E_{i,j}, 1 \leq i, j \leq n$ the canonical basis of $\mathfrak{gl}(n, \mathbb{C})$). The Cartan subalgebra $\mathfrak{h}$ is $\bigoplus_{i=1}^{i=n} \mathbb{C}H_i$ with $H_i = \tilde{E}_{i,i}$ and the nilradical of the Borel subalgebra is $D_n^+ = \bigoplus_{1 \leq i < j \leq n} \mathbb{C}\tilde{E}_{i,j} \oplus \bigoplus_{1 \leq i < j \leq n} \mathbb{C}\tilde{F}_{i,j}$. All $\tilde{F}_{i,j}$'s commute to one another, and one has:

$$[\tilde{E}_{i,j}, \tilde{F}_{k,l}] = \delta_{j,k} \tilde{F}_{i,l} - \delta_{j,l} \tilde{F}_{i,k} \quad (10)$$

We identify D_{n-1} to a subalgebra of D_n by simply taking the first row and first column of each block to be zero in (9). For $n=2$, $D_2^+ = \mathbb{C}^2$ is $\mathcal{I}$ -null. Suppose the result holds true for D_{n-1}^+ . One has

$$D_n^+ = \mathbb{C}\tilde{E}_{1,2} \oplus \mathbb{C}\tilde{E}_{1,3} \oplus \cdots \oplus \mathbb{C}\tilde{E}_{1,n} \oplus \mathbb{C}\tilde{F}_{1,n} \oplus \cdots \oplus \mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+. \quad (11)$$

Start with $\mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+$. From (10), $\tilde{F}_{1,2}$ commutes with all $\tilde{E}_{i,j}$ ($2 \leq i < j \leq n$) hence with D_{n-1}^+ . Then D_{n-1}^+ is a codimension 1 ideal of $\mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+$ and $\mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+$ is $\mathcal{I}$ -null from corr. 1. Consider now $\mathbb{C}\tilde{F}_{1,3} \oplus (\mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+)$. Again from (10), $\tilde{F}_{1,3}$ commutes with all elements of the basis of D_{n-1}^+ except $\tilde{E}_{2,3}$ and $[\tilde{E}_{2,3}, \tilde{F}_{1,3}] = \tilde{F}_{1,2}$. Then $\mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+$ is a codimension 1 ideal of $\mathbb{C}\tilde{F}_{1,3} \oplus (\mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+)$, and the latter is $\mathcal{I}$ -null. Suppose that $\mathbb{C}\tilde{F}_{1,s-1} \oplus \cdots \oplus \mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+$ is a codimension 1 ideal of $\mathbb{C}\tilde{F}_{1,s} \oplus (\mathbb{C}\tilde{F}_{1,s-1} \oplus \cdots \oplus \mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+)$, and that the latter is $\mathcal{I}$ -null. Consider $\mathbb{C}\tilde{F}_{1,s+1} \oplus (\mathbb{C}\tilde{F}_{1,s} \oplus \cdots \oplus \mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+)$. From (10), for $2 \leq i < j \leq n$, $[\tilde{E}_{i,j}, \tilde{F}_{1,s+1}] = \delta_{j,s+1} \tilde{F}_{1,i}$ is nonzero only for $i = 2, \dots, s$, and $j = s+1$, and it is then equal to $\tilde{F}_{1,i}$. Then first $\mathbb{C}\tilde{F}_{1,s} \oplus (\mathbb{C}\tilde{F}_{1,s-1} \oplus \cdots \oplus$

$\mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+$ is a codimension 1 ideal of $\mathbb{C}\tilde{F}_{1,s+1} \oplus (\mathbb{C}\tilde{F}_{1,s} \oplus \cdots \oplus \mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+)$. Second, the latter is $\mathcal{I}$ -null from corr. 1. By induction the above property holds for $s = n$. Consider now $\mathbb{C}\tilde{E}_{1,n} \oplus (\mathbb{C}\tilde{F}_{1,n} \oplus \cdots \oplus \mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+)$. One has for $2 \leq i < j \leq n$, $[\tilde{E}_{1,n}, \tilde{E}_{i,j}] = 0$, $[\tilde{E}_{1,n}, \tilde{F}_{i,j}] = -\delta_{n,j}\tilde{F}_{1,i}$, $[\tilde{E}_{1,n}, \tilde{F}_{1,j}] = 0$. Hence $\mathbb{C}\tilde{F}_{1,n} \oplus \cdots \oplus \mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+$ is an ideal of $\mathbb{C}\tilde{E}_{1,n} \oplus (\mathbb{C}\tilde{F}_{1,n} \oplus \cdots \oplus \mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+)$ and the latter is $\mathcal{I}$ -null. For $2 \leq i < j \leq n$, $1 \leq k \leq n-2$,

$$\begin{aligned} [\tilde{E}_{1,n-k}, \tilde{E}_{i,j}] &= \delta_{n-k,i}\tilde{E}_{1,j}, \\ [\tilde{E}_{1,n-k}, \tilde{F}_{i,j}] &= \delta_{n-k,i}\tilde{F}_{1,j} - \delta_{n-k,j}\tilde{F}_{1,i}, \\ [\tilde{E}_{1,n-k}, \tilde{E}_{1,n}] &= \delta_{n-k,1}\tilde{E}_{1,n}, \\ [\tilde{E}_{1,n-k}, \tilde{F}_{1,j}] &= \delta_{n-k,1}\tilde{F}_{1,j}. \end{aligned}$$

$[\tilde{E}_{1,n-1}, \tilde{E}_{i,j}]$ is nonzero only for $i = n-1, j = n$ and yields then $\tilde{E}_{1,n}$; $[\tilde{E}_{1,n-1}, \tilde{F}_{i,j}]$ is nonzero only for $(i = n-1, j = n)$ or for $(i < j = n-1)$ and yields respectively $\tilde{F}_{1,n}$, or $-\tilde{F}_{1,i}$. $[\tilde{E}_{1,n-1}, \tilde{E}_{1,n}]$ and $[\tilde{E}_{1,n-1}, \tilde{F}_{1,j}]$ are zero for $n \geq 3$. Hence, first $\mathbb{C}\tilde{E}_{1,n} \oplus \mathbb{C}\tilde{F}_{1,n} \oplus \cdots \oplus \mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+$ is a codimension 1 ideal of $\mathbb{C}\tilde{E}_{1,n-1} \oplus (\mathbb{C}\tilde{E}_{1,n} \oplus \mathbb{C}\tilde{F}_{1,n} \oplus \cdots \oplus \mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+)$, and second, the latter is $\mathcal{I}$ -null, since $\tilde{E}_{n-1,n}$ commutes with $\tilde{F}_{i,n-1}, \tilde{F}_{n-1,n}$. Suppose that $\mathbb{C}\tilde{E}_{1,n-k+1} \oplus \cdots \oplus \mathbb{C}\tilde{E}_{1,n} \oplus \mathbb{C}\tilde{F}_{1,n} \oplus \cdots \oplus \mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+$ is a codimension 1 ideal of $\mathbb{C}\tilde{E}_{1,n-k} \oplus (\mathbb{C}\tilde{E}_{1,n-k+1} \oplus \cdots \oplus \mathbb{C}\tilde{E}_{1,n} \oplus \mathbb{C}\tilde{F}_{1,n} \oplus \cdots \oplus \mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+)$ and that the latter is $\mathcal{I}$ -null. Consider $\mathbb{C}\tilde{E}_{1,n-k-1} \oplus (\mathbb{C}\tilde{E}_{1,n-k} \oplus \cdots \oplus \mathbb{C}\tilde{E}_{1,n} \oplus \mathbb{C}\tilde{F}_{1,n} \oplus \cdots \oplus \mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+)$. $[\tilde{E}_{1,n-k-1}, \tilde{E}_{i,j}]$ is nonzero only for $i = n-k-1$ and yields then $\tilde{E}_{1,j}$; $[\tilde{E}_{1,n-k-1}, \tilde{F}_{i,j}] = \delta_{n-k-1,i}\tilde{F}_{1,j} - \delta_{n-k-1,j}\tilde{F}_{1,i}$ is nonzero only for $i = n-k-1$ or $j = n-k-1$ and yields resp. $\tilde{F}_{1,j}$ or $-\tilde{F}_{1,i}$. Hence $\mathbb{C}\tilde{E}_{1,n-k} \oplus \cdots \oplus \mathbb{C}\tilde{E}_{1,n} \oplus \mathbb{C}\tilde{F}_{1,n} \oplus \cdots \oplus \mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+$ is an ideal of $\mathbb{C}\tilde{E}_{1,n-k-1} \oplus (\mathbb{C}\tilde{E}_{1,n-k} \oplus \cdots \oplus \mathbb{C}\tilde{E}_{1,n} \oplus \mathbb{C}\tilde{F}_{1,n} \oplus \cdots \oplus \mathbb{C}\tilde{F}_{1,2} \oplus D_{n-1}^+)$. The latter is $\mathcal{I}$ -null since $\tilde{E}_{n-k-1,j}$ commutes with both $\tilde{F}_{n-k-1,j'}$, $\tilde{F}_{i,n-k-1}$. The result follows by induction.

Case B_n . We may take B_n ($n \geq 2$) as the Lie algebra of matrices

$$\left(\begin{array}{c|cc} 0 & u & v \\ \hline -{}^t v & Z_1 & Z_2 \\ \hline -{}^t u & Z_3 & -{}^t Z_1 \end{array} \right) \quad (12)$$

with u, v complex $(1 \times n)$ -matrices, $Z_i \in \mathfrak{gl}(n, \mathbb{C})$, Z_2, Z_3 skew symmetric, i.e.

$$\left(\begin{array}{c|cc} 0 & u & v \\ \hline -{}^t v & & \\ \hline -{}^t u & \boxed{A} & \end{array} \right) \quad (13)$$

with $A \in D_n$. The Cartan subalgebra is the same as for D_n . Then B_n^+ consists of the matrices

$$\left(\begin{array}{c|cc} 0 & 0 & v \\ \hline -{}^t v & & \\ \hline 0 & \boxed{A} & \end{array} \right) \quad (14)$$

with v complex $(1 \times n)$ -matrix and $A \in D_n^+$. We identify $A \in D_n^+$ to the matrix

$$\left(\begin{array}{c|cc} 0 & 0 & 0 \\ \hline 0 & & \\ \hline & \boxed{A} & \\ \hline 0 & & \end{array} \right) \in B_n^+.$$

For $1 \leq q \leq n$, let v_q the $1 \times n$ -matrix $E_{1,q} = (0, \dots, 1, \dots, 0)$ (1 in q^{th} position), and

$$\tilde{v}_q = \left(\begin{array}{c|cc} 0 & 0 & v_q \\ \hline -{}^t v_q & & \\ \hline & \boxed{0} & \\ \hline 0 & & \end{array} \right)$$

Hence $B_n^+ = \left(\bigoplus_{q=1}^n \mathbb{C} \tilde{v}_q \right) \oplus D_n^+$. One has for $1 \leq q \leq n$, $1 \leq i < j \leq n$

$$\begin{aligned} [\tilde{v}_q, \tilde{E}_{i,j}] &= -\delta_{q,j} \tilde{v}_i \\ [\tilde{v}_q, \tilde{F}_{i,j}] &= 0 \end{aligned}$$

and for $1 \leq s < q \leq n$

$$[\tilde{v}_q, \tilde{v}_s] = \tilde{F}_{s,q}. \quad (15)$$

Consider $\mathbb{C} \tilde{v}_1 \oplus D_n^+$. As $\tilde{v}_1$ commutes with $\tilde{E}_{i,j}, \tilde{F}_{i,j}$, D_n^+ is an ideal of $\mathbb{C} \tilde{v}_1 \oplus D_n^+$ and the latter is $\mathcal{I}$ -null. Suppose $\mathbb{C} \tilde{v}_{s-1} \oplus \dots \oplus \mathbb{C} \tilde{v}_1 \oplus D_n^+$ is an ideal of $\mathbb{C} \tilde{v}_s \oplus (\mathbb{C} \tilde{v}_{s-1} \oplus \dots \oplus \mathbb{C} \tilde{v}_1 \oplus D_n^+)$ and the latter is $\mathcal{I}$ -null. Consider $\mathbb{C} \tilde{v}_{s+1} \oplus (\mathbb{C} \tilde{v}_s \oplus \mathbb{C} \tilde{v}_{s-1} \oplus \dots \oplus \mathbb{C} \tilde{v}_1 \oplus D_n^+)$. $[\tilde{v}_{s+1}, \tilde{E}_{i,j}] = -\delta_{s+1,j} \tilde{v}_i$ hence $\tilde{v}_{s+1}$ commutes to all $\tilde{E}_{i,j}$ s except for $\tilde{E}_{i,s+1}$ ($i < s+1$) and then yields $-\tilde{v}_i$. For $t < s+1$, $[\tilde{v}_{s+1}, \tilde{v}_t] = \tilde{F}_{t,s+1}$. Hence $\mathbb{C} \tilde{v}_s \oplus \mathbb{C} \tilde{v}_{s-1} \oplus \dots \oplus \mathbb{C} \tilde{v}_1 \oplus D_n^+$ is an ideal of $\mathbb{C} \tilde{v}_{s+1} \oplus (\mathbb{C} \tilde{v}_s \oplus \mathbb{C} \tilde{v}_{s-1} \oplus \dots \oplus \mathbb{C} \tilde{v}_1 \oplus D_n^+)$. The latter is $\mathcal{I}$ -null since the $\tilde{E}_{i,s+1}$ ($i < s+1$) commute to one another. Hence by induction the property holds for $s = n$ and B_n^+ is $\mathcal{I}$ -null.

Case C_n . This case is pretty similar to the case D_n . We may take C_n as the Lie algebra of matrices

$$\begin{pmatrix} Z_1 & Z_2 \\ Z_3 & -{}^t Z_1 \end{pmatrix} \quad (16)$$

with $Z_i \in \mathfrak{gl}(n, \mathbb{C})$, Z_2, Z_3 symmetric. $\tilde{E}_{i,j}$ and the Cartan subalgebra are identical to those of D_n . We denote for $1 \leq i, j \leq n$: $\hat{F}_{i,j} = \begin{pmatrix} 0 & E_{i,j} + E_{j,i} \\ 0 & 0 \end{pmatrix}$. Then

$$C_n^+ = \bigoplus_{1 \leq i < j \leq n} \mathbb{C} \tilde{E}_{i,j} \oplus \bigoplus_{1 \leq k \leq l \leq n} \mathbb{C} \hat{F}_{k,l}. \quad (17)$$

All $\hat{F}_{k,l}$'s commute to one another, and one has:

$$[\tilde{E}_{i,j}, \hat{F}_{k,l}] = \delta_{j,k} \hat{F}_{i,l} + \delta_{j,l} \hat{F}_{i,k} \quad (18)$$

The case is step by step analog to the case of D_n with (18) instead of (10) and (17) instead of (11).

Case G_2 . The commutation relations for G_2 appear in [6], p. 346. G_2^+ is 6-dimensional with comm. rel. $[x_1, x_2] = x_3$; $[x_1, x_3] = x_4$; $[x_1, x_4] = -3x_5$; $[x_2, x_5] = -x_6$; $[x_3, x_4] = -3x_6$. G_2^+ has the same adjoint cohomology $(1, 4, 7, 8, 7, 5, 2)$ as, and is isomorphic to, $\mathfrak{g}_{6,18}$, which is $\mathcal{I}$ -null.

Case F_4 . F_4^+ has 24 positive roots, and root vectors x_i ($1 \leq i \leq 24$). From the root pattern, one gets with some calculations the commutation relations of F_4^+ : $[x_1, x_2] = x_5$; $[x_1, x_{13}] = x_{14}$; $[x_1, x_{15}] = -x_6$; $[x_1, x_{16}] = -x_7$; $[x_1, x_{17}] = -x_{23}$; $[x_1, x_{18}] = x_{19}$; $[x_1, x_{24}] = x_{22}$; $[x_2, x_3] = x_{15}$; $[x_2, x_7] = x_8$; $[x_2, x_{12}] = x_{13}$; $[x_2, x_{19}] = x_{20}$; $[x_2, x_{21}] = x_{24}$; $[x_2, x_{23}] = x_9$; $[x_3, x_4] = x_{21}$; $[x_3, x_5] = x_6$; $[x_3, x_6] = x_7$; $[x_3, x_9] = x_{10}$; $[x_3, x_{11}] = x_{12}$; $[x_3, x_{15}] = x_{16}$; $[x_3, x_{20}] = -2x_{11}$; $[x_3, x_{22}] = \frac{1}{2}x_{23}$; $[x_3, x_{24}] = -\frac{1}{2}x_{17}$; $[x_4, x_6] = x_{22}$; $[x_4, x_7] = x_{23}$; $[x_4, x_8] = x_9$; $[x_4, x_9] = -x_{20}$; $[x_4, x_{10}] = x_{11}$; $[x_4, x_{15}] = -x_{24}$; $[x_4, x_{16}] = x_{17}$; $[x_4, x_{17}] = x_{18}$; $[x_4, x_{23}] = -x_{19}$; $[x_5, x_{12}] = x_{14}$; $[x_5, x_{16}] = x_8$; $[x_5, x_{17}] = x_9$; $[x_5, x_{18}] = -x_{20}$; $[x_5, x_{21}] = x_{22}$; $[x_6, x_{11}] = -x_{14}$; $[x_6, x_{15}] = -x_8$; $[x_6, x_{17}] = x_{10}$; $[x_6, x_{18}] = 2x_{11}$; $[x_6, x_{21}] = \frac{1}{2}x_{23}$; $[x_6, x_{24}] = \frac{1}{2}x_9$; $[x_7, x_{18}] = 2x_{12}$; $[x_7, x_{20}] = -2x_{14}$; $[x_7, x_{24}] = x_{10}$; $[x_8, x_{18}] = 2x_{13}$; $[x_8, x_{19}] = 2x_{14}$; $[x_8, x_{21}] = -x_{10}$; $[x_9, x_{17}] = -2x_{13}$; $[x_9, x_{21}] = -x_{11}$; $[x_9, x_{23}] = 2x_{14}$; $[x_{10}, x_{21}] = -x_{12}$; $[x_{10}, x_{22}] = -x_{14}$; $[x_{10}, x_{24}] = -x_{13}$; $[x_{11}, x_{15}] = -x_{13}$; $[x_{15}, x_{19}] = 2x_{11}$; $[x_{15}, x_{21}] = \frac{1}{2}x_{17}$; $[x_{15}, x_{22}] = \frac{1}{2}x_9$; $[x_{15}, x_{23}] = -x_{10}$; $[x_{16}, x_{19}] = 2x_{12}$; $[x_{16}, x_{20}] = 2x_{13}$; $[x_{16}, x_{22}] = x_{10}$; $[x_{17}, x_{22}] = x_{11}$; $[x_{17}, x_{23}] = 2x_{12}$; $[x_{21}, x_{22}] = \frac{1}{2}x_{19}$; $[x_{21}, x_{24}] = \frac{1}{2}x_{18}$; $[x_{22}, x_{24}] = -\frac{1}{2}x_{20}$; $[x_{23}, x_{24}] = x_{11}$. Then one verifies (see [16]) that F_4^+ is $\mathcal{I}$ -null.

Case E_6 . In the case of E_6^+ the set Δ_+ of positive roots (associated to the set S of simple roots) has cardinality 36 ([6], p. 333):

$$\begin{aligned} \Delta_+ = & \{\varepsilon_i + \varepsilon_j; 1 \leq i < j \leq 5\} \cup \{\varepsilon_i - \varepsilon_j; 1 \leq j < i \leq 5\} \\ & \cup \left\{ \frac{1}{2}(\pm\varepsilon_1 \pm \varepsilon_2 \pm \varepsilon_3 \pm \varepsilon_4 \pm \varepsilon_5 + \sqrt{3}\varepsilon_6); \# \text{ minus signs even} \right\}. \end{aligned}$$

Instead of computing commutation relations, we will use the following property of Δ_+ .

($\mathcal{P}$): for $\alpha, \beta, \gamma \in \Delta_+$, if $\alpha + \beta \in \Delta_+$ and $\alpha + \gamma \in \Delta_+$ then $\beta + \gamma \notin \Delta_+$.

Now, introduce some Chevalley basis ([7], p. 19 ex. 7) of $E_6^+ : (X_\alpha)_{\alpha \in \Delta_+}$. One has

$$[X_\alpha, X_\beta] = N_{\alpha, \beta} X_{\alpha + \beta} \quad \forall \alpha, \beta \in \Delta_+$$

$$N_{\alpha, \beta} = 0 \text{ if } \alpha + \beta \notin \Delta_+, N_{\alpha, \beta} \in \mathbb{Z} \setminus \{0\} \text{ if } \alpha + \beta \in \Delta_+.$$

Define inductively a sequence $\mathfrak{g}_1 \subset \mathfrak{g}_2 \subset \dots \subset \mathfrak{g}_{36} = E_6^+$ of $\mathcal{I}$ -null subalgebras, each of which a codimension 1 ideal of the following, as follows. Start with $\mathfrak{g}_1 = \mathbb{C}X_{\delta_1}$, $\delta_1 \in \Delta_+$ of maximum height. Suppose $\mathfrak{g}_i$ defined. Then take $\mathfrak{g}_{i+1} = \mathbb{C}X_{\delta_{i+1}} \oplus \mathfrak{g}_i$ with $\delta_{i+1} \in \Delta_+ \setminus \{\delta_1, \dots, \delta_i\}$ of maximum height. Clearly, $\mathfrak{g}_i$ is a 1-dimensional ideal of $\mathfrak{g}_{i+1}$. To prove that it is $\mathcal{I}$ -null we only have to check that, for $1 \leq s, t \leq i$, if $\delta_{i+1} + \delta_s \in \Delta_+$ and $\delta_{i+1} + \delta_t \in \Delta_+$ then $\delta_s + \delta_t \notin \Delta_+$. That holds true because of property ($\mathcal{P}$).

Case E_7 . In the case of E_7^+ the set Δ_+ of positive roots (associated to the

set S of simple roots) has cardinality 63 ([6], p. 333):

$$\begin{aligned}\Delta_+ &= \{\varepsilon_i + \varepsilon_j; 1 \leq i < j \leq 6\} \cup \{\varepsilon_i - \varepsilon_j; 1 \leq j < i \leq 6\} \cup \{\sqrt{2}\varepsilon_7\} \\ &\cup \left\{ \frac{1}{2}(\pm\varepsilon_1 \pm \varepsilon_2 \pm \varepsilon_3 \pm \varepsilon_4 \pm \varepsilon_5 \pm \varepsilon_6 + \sqrt{2}\varepsilon_7); \# \text{ minus signs odd} \right\}.\end{aligned}$$

Property $(\mathcal{P})$ holds true for E_7^+ (see [16]). Hence the conclusion follows as in the case of E_6^+ .

Case E_8 . In the case of E_8^+ the set Δ_+ of positive roots has cardinality 120 ([6], p. 333):

$$\begin{aligned}\Delta_+ &= \{\varepsilon_i + \varepsilon_j; 1 \leq i < j \leq 8\} \cup \{\varepsilon_i - \varepsilon_j; 1 \leq j < i \leq 8\} \\ &\cup \left\{ \frac{1}{2}(\pm\varepsilon_1 \pm \varepsilon_2 \pm \varepsilon_3 \pm \varepsilon_4 \pm \varepsilon_5 \pm \varepsilon_6 \pm \varepsilon_7 + \varepsilon_8); \# \text{ minus signs even} \right\}.\end{aligned}$$

Property $(\mathcal{P})$ holds true for E_8^+ (see [16]). Hence the conclusion follows as in the case of E_6^+ . $\square$

Remark 1. Property $(\mathcal{P})$ holds true for A_n^+ , hence we could have used it. However, it does not hold true for F_4^+ . One has for example in the above commutation relations of F_4^+ (with root vectors) $[x_3, x_4] \neq 0$, $[x_3, x_9] \neq 0$, yet $[x_4, x_9] \neq 0$.

Corollary 5. Suppose that the Lie algebra $\mathfrak{g}$ is such that $\dim \text{Im } \mathcal{I} = 0$ or 1. Let $\tau \in \text{Der } \mathfrak{g}$ such that $\tau x_k \in \mathcal{C}^2 \mathfrak{g} \forall k \geq 2$ where $(x_1, \dots, x_N)$ is some basis of $\mathfrak{g}$. Denote $\tilde{\mathfrak{g}}_\tau = \mathbb{C}\tau \oplus \mathfrak{g}$ the Lie algebra obtained by adjoining the derivation τ to $\mathfrak{g}$, and by $\tilde{\mathcal{I}}$ the Koszul map of $\tilde{\mathfrak{g}}_\tau$. Then $\dim \text{Im } \tilde{\mathcal{I}} = 0$ or 1.

Proof. Let $B \in (S^2 \tilde{\mathfrak{g}}_\tau^*)^{\tilde{\mathfrak{g}}_\tau}$ such that $I_B \neq 0$. One has $I_B = \omega^\tau \wedge df_\tau + I_{B_2} \circ (\pi_2 \times \pi_2 \times \pi_2)$ where $(\tau, x_1, \dots, x_N)$ is the basis of $\tilde{\mathfrak{g}}_\tau$, $(\omega^\tau, \omega^1, \dots, \omega^N)$ the dual basis, B_2 the restriction of B to $\mathfrak{g}$, $f_\tau = B(\tau, \cdot)$ and π_2 the projection on $\mathfrak{g}$. One may suppose $\dim \text{Im } \mathcal{I} = 1$. Let $C \in (S^2 \mathfrak{g}^*)^{\mathfrak{g}}$ with $I_C \neq 0$. There exists $\lambda \in \mathbb{C}$ such that $I_{B_2} = \lambda I_C$. Now, for $X, Y \in \mathfrak{g}$,

$$\begin{aligned}df_\tau(X, Y) &= B(\tau, [X, Y]) \\ &= \omega^1(X)B(\tau, [\pi_3(Y)]) - \omega^1(Y)B(\tau, [\pi_3(X)]) + B(\tau, [\pi_3(X), \pi_3(Y)]) \\ &= -\omega^1(X)B(\tau\pi_3(Y), x_1) + \omega^1(Y)B(\tau\pi_3(X), x_1) + B(\tau\pi_3(X), \pi_3(Y))\end{aligned}$$

where π_3 is the projection on $\text{vect}(x_2, \dots, x_N)$. Now, $B(\tau\pi_3(Y), x_1) = \lambda C(\tau\pi_3(Y), x_1)$, $B(\tau\pi_3(X), x_1) = \lambda C(\tau\pi_3(X), x_1)$, $B(\tau\pi_3(X), \pi_3(Y)) = \lambda C(\tau\pi_3(X), \pi_3(Y))$ since $\tau\pi_3(X), \tau\pi_3(Y) \in \mathcal{C}^2 \mathfrak{g}$. The result follows. $\square$

Example 2. The nilpotent Lie algebra $\mathfrak{g}_{7,2,4}$, has commutation relations $[x_1, x_2] = x_3$, $[x_1, x_3] = x_4$, $[x_1, x_4] = x_5$, $[x_1, x_5] = x_6$, $[x_2, x_5] = -x_7$, $[x_3, x_4] = x_7$. The

elements of $\text{Der } \mathfrak{g}_{7,2.4}(\text{mod } \text{ad}_{\mathfrak{g}_{7,2.4}})$ are

$$\tau = \begin{pmatrix} \xi_1^1 & 0 & 0 & 0 & 0 & 0 & 0 \\ \xi_1^2 & \xi_2^2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \xi_1^1 + \xi_2^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2\xi_1^1 + \xi_2^2 & 0 & 0 & 0 \\ 0 & \xi_2^5 & 0 & 0 & 3\xi_1^1 + \xi_2^2 & 0 & 0 \\ \xi_1^6 & \xi_2^6 & \xi_2^5 & 0 & 0 & 4\xi_1^1 + \xi_2^2 & 0 \\ \xi_1^7 & 0 & 0 & 0 & 0 & -\xi_2^1 & 3\xi_1^1 + 2\xi_2^2 \end{pmatrix} \quad (19)$$

τ is nilpotent if $\xi_1^1 = \xi_2^2 = 0$. Denote the nilpotent τ by $(\xi_1^2; \xi_2^5; \xi_1^6, \xi_2^6; \xi_1^7)$. Now, projectively equivalent derivations τ, τ' (see [14]) give isomorphic $\tilde{\mathfrak{g}}_\tau, \tilde{\mathfrak{g}}_{\tau'}$. By reduction using projective equivalence, we are reduced to the following cases: Case 1. $\xi_1^2 \neq 0 : (1; \varepsilon; 0, \eta; 0)$; Case 2. $\xi_1^2 = 0 : (0; \varepsilon; 0, \eta; \lambda)$; where $\varepsilon, \eta, \lambda = 0, 1$. In both cases $\tilde{\mathfrak{g}}_\tau$ is $\mathcal{I}$ -null, except when $\tau = 0$ in case 2 where $\tilde{\mathfrak{g}}_\tau$ is the direct product $\mathbb{C} \times \mathfrak{g}_{7,2.4}$ which is quadratic (i.e. having a nondegenerate invariant bilinear form). Hence any indecomposable 8-dimensional nilpotent Lie algebra containing a subalgebra isomorphic to $\mathfrak{g}_{7,2.4}$ is $\mathcal{I}$ -null, though $\mathfrak{g}_{7,2.4}$ is quadratic. That is in line with the fact that, from the double extension method of [20], [17], any indecomposable quadratic solvable Lie algebra is a *double extension* of a quadratic solvable Lie algebra by $\mathbb{C}$.

Example 3. Among the 170 (non isomorphic) nilpotent complex Lie algebras of dimension ≤ 7 , only a few are not $\mathcal{I}$ -null. Those are listed in Table 1 in the classification of [13], [15] (they are all $\mathcal{I}$ -exact). Table 1 gives for each of them $\dim (S^2 \mathfrak{g}^*)^{\mathfrak{g}}$, a basis for $((S^2 \mathfrak{g}^*)^{\mathfrak{g}} / \ker \mathcal{I})$ (which in those cases is one-dimensional), and the corresponding I_{Bs} . ■ denotes quadratic Lie algebras; for $\omega, \pi \in \mathfrak{g}^*$, $\circ$ stands for the symmetric product $\omega \circ \pi = \omega \otimes \pi + \pi \otimes \omega$; $\omega^{i,j,k}$ stands for $\omega^i \wedge \omega^j \wedge \omega^k$.

Remark 2. There are nilpotent Lie algebras of higher dimension with $\dim (S^2 \mathfrak{g}^*)^{\mathfrak{g}} / \ker \mathcal{I} > 1$. For example, in the case of the 10 dimensional Lie algebra $\mathfrak{g}$ with commutation relations $[x_1, x_2] = x_5, [x_1, x_3] = x_6, [x_1, x_4] = x_7, [x_2, x_3] = x_8, [x_2, x_4] = x_9, [x_3, x_4] = x_{10}$, $\dim (S^2 \mathfrak{g}^*)^{\mathfrak{g}} / \ker \mathcal{I} = 4$, and in the analogous case of the 15 dimensional nilpotent Lie algebra with 5 generators one has $\dim (S^2 \mathfrak{g}^*)^{\mathfrak{g}} / \ker \mathcal{I} = 10$. Those algebras are $\mathcal{I}$ -exact and not quadratic.

Remark 3. In the *transversal to dimension* approach to the classification problem of nilpotent Lie algebras initiated in [21], one first associates a generalized Cartan matrix (abbr. GCM) A to any nilpotent finite dimensional complex Lie algebra $\mathfrak{g}$, and then looks at $\mathfrak{g}$ as the quotient $\hat{\mathfrak{g}}(A)_+ / \mathcal{J}$ of the nilradical of the Borel subalgebra of the Kac-Moody Lie algebra $\hat{\mathfrak{g}}(A)$ associated to A by some ideal $\mathcal{J}$. Then one gets for any GCM A the subproblem of classifying (up to the action of a certain group) all ideals of $\hat{\mathfrak{g}}(A)_+$, thus getting all nilpotent Lie algebras of type A (see [1], [2], [3], [22], and the references therein). Any indecomposable GCM is of exactly one of the 3 types *finite*, *affine*, *indefinite*

Table 1:

algebra	$\dim (S^2 \mathfrak{g}^*)^{\mathfrak{g}}$	basis for $(S^2 \mathfrak{g}^*)^{\mathfrak{g}} / \ker \mathcal{I}$	I_B .
$\mathfrak{g}_{5,4}$ ■	4	$\omega^1 \odot \omega^5 - \omega^2 \odot \omega^4 + \omega^3 \otimes \omega^3$	$\omega^{1,2,3} = d\omega^{1,5}$
$\mathfrak{g}_{6,3}$ ■	7	$\omega^1 \odot \omega^6 - \omega^2 \odot \omega^5 + \omega^3 \odot \omega^4$	$\omega^{1,2,3} = d\omega^{1,6}$
$\mathfrak{g}_{6,14}$	4	$\omega^1 \odot \omega^6 - \omega^2 \odot \omega^4 + \omega^3 \otimes \omega^3$	$\omega^{1,2,3} = -d\omega^{1,4}$
$\mathfrak{g}_{5,4} \times \mathbb{C}$ ■	7	$\omega^1 \odot \omega^5 - \omega^2 \odot \omega^4 + \omega^3 \otimes \omega^3$	$\omega^{1,2,3} = d\omega^{1,5}$
$\mathfrak{g}_{5,4} \times \mathbb{C}^2$ ■	11	$\omega^1 \odot \omega^5 - \omega^2 \odot \omega^4 + \omega^3 \otimes \omega^3$	$\omega^{1,2,3} = d\omega^{1,5}$
$\mathfrak{g}_{6,3} \times \mathbb{C}$ ■	11	$\omega^1 \odot \omega^6 - \omega^2 \odot \omega^5 + \omega^3 \odot \omega^4$	$\omega^{1,2,3} = d\omega^{1,6}$
$\mathfrak{g}_{7,0.4}(\lambda),$ $\mathfrak{g}_{7,0.5},$ $\mathfrak{g}_{7,0.6},$ $\mathfrak{g}_{7,1.02},$ $\mathfrak{g}_{7,1.10},$ $\mathfrak{g}_{7,1.13},$ $\mathfrak{g}_{7,1.14},$ $\mathfrak{g}_{7,1.17}$	4	$\omega^1 \odot \omega^5 - \omega^2 \odot \omega^4 + \omega^3 \otimes \omega^3$	$\omega^{1,2,3} = d\omega^{1,5}$
$\mathfrak{g}_{7,1.03}$	4	$\omega^1 \odot \omega^6 - \omega^2 \odot \omega^4 + \omega^3 \otimes \omega^3$	$\omega^{1,2,3} = d\omega^{1,6}$
$\mathfrak{g}_{7,2.2}$	7	$\omega^1 \odot \omega^4 - \omega^2 \odot \omega^6 + \omega^3 \odot \omega^5$	$\omega^{1,2,3} = d\omega^{1,4}$
$\mathfrak{g}_{7,2.4}$ ■	4	$\omega^1 \odot \omega^7 + \omega^2 \odot \omega^6 - \omega^3 \odot \omega^5 + \omega^4 \otimes \omega^4$	$\omega^{1,3,4} - \omega^{1,2,5} = d\omega^{1,7}$
$\mathfrak{g}_{7,2.5},$ $\mathfrak{g}_{7,2.6},$ $\mathfrak{g}_{7,2.7},$ $\mathfrak{g}_{7,2.8},$ $\mathfrak{g}_{7,2.9},$	4	$\omega^1 \odot \omega^5 - \omega^2 \odot \omega^4 + \omega^3 \otimes \omega^3$	$\omega^{1,2,3} = d\omega^{1,5}$
$\mathfrak{g}_{7,2.18}$	7	$\omega^1 \odot \omega^6 - \omega^2 \odot \omega^5 + \omega^4 \otimes \omega^4$	$\omega^{1,2,4} = d\omega^{1,6}$
$\mathfrak{g}_{7,2.44},$ $\mathfrak{g}_{7,3.6}$	7	$\omega^1 \odot \omega^6 - \omega^2 \odot \omega^5 + \omega^3 \odot \omega^4$	$\omega^{1,2,3} = d\omega^{1,6}$
$\mathfrak{g}_{7,3.23}$	7	$\omega^1 \odot \omega^6 - \omega^2 \odot \omega^5 + \omega^3 \otimes \omega^3$	$\omega^{1,2,3} = d\omega^{1,6}$

(among that last the hyperbolic GCMs, with the property that any connected proper subdiagram of the Dynkin diagram is of finite or affine type) ([8],[24]). From corollary 4, the nilpotent Lie algebras that are not $\mathcal{I}$ -null all come from affine or indefinite types. Unfortunately, that is the case of many nilpotent Lie algebras, see table 2.

Example 4. The quadratic 5-dimensional nilpotent Lie algebra $\mathfrak{g}_{5,4}$ has commutation relations $[x_1, x_2] = x_3$, $[x_1, x_3] = x_4$, $[x_2, x_3] = x_5$. Consider the 10-dimensional direct product $\mathfrak{g}_{5,4} \times \mathfrak{g}_{5,4}$, with the commutation relations: $[x_1, x_2] = x_5$, $[x_1, x_5] = x_6$, $[x_2, x_5] = x_7$, $[x_3, x_4] = x_8$, $[x_3, x_8] = x_9$, $[x_4, x_8] = x_{10}$. The only 11-dimensional nilpotent Lie algebra with an invariant bilinear form which reduces to $B_1 = \omega^1 \odot \omega^7 - \omega^2 \odot \omega^6 + \omega^5 \otimes \omega^5$, $B_2 = \omega^3 \odot \omega^{10} - \omega^4 \odot \omega^9 + \omega^8 \otimes \omega^8$, on respectively the first and second factor is the direct product $\mathbb{C} \times \mathfrak{g}_{5,4} \times \mathfrak{g}_{5,4}$,

Example 5. The 4-dimensional solvable "diamond" Lie algebra $\mathfrak{g}$ with basis (x_1, x_2, x_3, x_4) and commutation relations $[x_1, x_2] = x_3$, $[x_1, x_3] = -x_2$, $[x_2, x_3] = x_4$ cannot be obtained as in lemma 3. Here $\dim \left((S^2 \mathfrak{g}^*)^{\mathfrak{g}} / \ker \mathcal{I} \right) = 1$, with basis element $B = \omega^1 \odot \omega^4 + \omega^2 \otimes \omega^2 + \omega^3 \otimes \omega^3$. $I_B = \omega^{1,2,3} = d\omega^{1,4}$; $\mathfrak{g}$ is quadratic and $\mathcal{I}$ -exact. In fact, one verifies that all other solvable 4-dimensional Lie solvable Lie algebras are $\mathcal{I}$ -null (for a list, see e.g. [18]).

4 Comparison of HL^2 and H^2 for a Lie algebra.

If $\mathfrak{c}$ denotes the center of $\mathfrak{g}$, $\mathfrak{c} \otimes (S^2 \mathfrak{g}^*)^{\mathfrak{g}}$ is the space of invariant $\mathfrak{c}$ -valued symmetric bilinear map and we denote $F = Id \otimes \mathcal{I} : \mathfrak{c} \otimes (S^2 \mathfrak{g}^*)^{\mathfrak{g}} \rightarrow C^3(\mathfrak{g}, \mathfrak{g}) = \mathfrak{g} \otimes \bigwedge^3 \mathfrak{g}^*$. Then $\text{Im } F = \mathfrak{c} \otimes \text{Im } \mathcal{I}$.

Theorem 1. *Let $\mathfrak{g}$ be any finite dimensional complex Lie algebra and $ZL_0^2(\mathfrak{g}, \mathfrak{g})$ (resp. $ZL_0^2(\mathfrak{g}, \mathbb{C})$) the space of symmetric adjoint (resp. trivial) Leibniz 2-cocycles.*

- (i) $ZL^2(\mathfrak{g}, \mathfrak{g}) / (Z^2(\mathfrak{g}, \mathfrak{g}) \oplus ZL_0^2(\mathfrak{g}, \mathfrak{g})) \cong (\mathfrak{c} \otimes \text{Im } \mathcal{I}) \cap B^3(\mathfrak{g}, \mathfrak{g})$.
- (ii) $ZL_0^2(\mathfrak{g}, \mathfrak{g}) = \mathfrak{c} \otimes \ker \mathcal{I}$. In particular, $\dim ZL_0^2(\mathfrak{g}, \mathfrak{g}) = c \frac{p(p+1)}{2}$ where $c = \dim \mathfrak{c}$ and $p = \dim \mathfrak{g} / C^2 \mathfrak{g} = \dim H^1(\mathfrak{g}, \mathbb{C})$.
- (iii) $HL^2(\mathfrak{g}, \mathfrak{g}) \cong H^2(\mathfrak{g}, \mathfrak{g}) \oplus (\mathfrak{c} \otimes \ker \mathcal{I}) \oplus ((\mathfrak{c} \otimes \text{Im } \mathcal{I}) \cap B^3(\mathfrak{g}, \mathfrak{g}))$.
- (iv) $ZL^2(\mathfrak{g}, \mathbb{C}) / (Z^2(\mathfrak{g}, \mathbb{C}) \oplus ZL_0^2(\mathfrak{g}, \mathbb{C})) \cong \text{Im } \mathcal{I} \cap B^3(\mathfrak{g}, \mathbb{C})$.
- (v) $ZL_0^2(\mathfrak{g}, \mathbb{C}) = \ker \mathcal{I}$.
- (vi) $HL^2(\mathfrak{g}, \mathbb{C}) \cong H^2(\mathfrak{g}, \mathbb{C}) \oplus \ker \mathcal{I} \oplus (\text{Im } \mathcal{I} \cap B^3(\mathfrak{g}, \mathbb{C}))$.

Proof. (i) The Leibniz 2-cochain space $CL^2(\mathfrak{g}, \mathfrak{g}) = \mathfrak{g} \otimes (\mathfrak{g}^*)^{\otimes 2}$ decomposes as $(\mathfrak{g} \otimes \bigwedge^2 \mathfrak{g}^*) \oplus (\mathfrak{g} \otimes S^2 \mathfrak{g}^*)$ with $\mathfrak{g} \otimes S^2 \mathfrak{g}^*$ the space of symmetric elements in $CL^2(\mathfrak{g}, \mathfrak{g})$. By definition of the Leibniz coboundary δ , one has for $\psi \in CL^2(\mathfrak{g}, \mathfrak{g})$ and $X, Y, Z \in \mathfrak{g}$

$$(\delta\psi)(X, Y, Z) = u + v + w + r + s + t \quad (20)$$

with $u = [X, \psi(Y, Z)]$, $v = [\psi(X, Z), Y]$, $w = -[\psi(X, Y), Z]$, $r = -\psi([X, Y], Z)$, $s = \psi(X, [Y, Z])$, $t = \psi([X, Z], Y)$. δ coincides with the usual coboundary operator on $\mathfrak{g} \otimes \bigwedge^2 \mathfrak{g}^*$. Now, let $\psi = \psi_1 + \psi_0 \in CL^2(\mathfrak{g}, \mathfrak{g})$, $\psi_1 \in \mathfrak{g} \otimes \bigwedge^2 \mathfrak{g}^*$, $\psi_0 \in \mathfrak{g} \otimes S^2 \mathfrak{g}^*$.

Suppose $\psi \in ZL^2(\mathfrak{g}, \mathfrak{g}) : \delta\psi = 0 = \delta\psi_1 + \delta\psi_0 = d\psi_1 + \delta\psi_0$. Then $\delta\psi_0 = -d\psi_1 \in \mathfrak{g} \otimes \bigwedge^3 \mathfrak{g}^*$ is antisymmetric. Then permuting X and Y in formula (20) for ψ_0 yields $(\delta\psi_0)(Y, X, Z) = -v - u + w - r + t + s$. As $\delta\psi_0$ is antisymmetric, we get

$$w + s + t = 0. \quad (21)$$

Now, the circular permutation (X, Y, Z) in (20) for ψ_0 yields $(\delta\psi_0)(Y, Z, X) = -v - w + u - s - t + r$. Again, by antisymmetry, we get

$$v + w + s + t = 0, \quad (22)$$

i.e. $(\delta\psi_0)(X, Y, Z) = u + r$. From (21) and (22), $v = 0$. Applying twice the circular permutation (X, Y, Z) to v , we get first $w = 0$ and then $u = 0$. Hence $(\delta\psi_0)(X, Y, Z) = r = -\psi_0([X, Y], Z)$. Note first that $u = 0$ reads $[X, \psi_0(Y, Z)] = 0$. As X, Y, Z are arbitrary, ψ_0 is $\mathfrak{c}$ -valued. Now the permutation of Y and Z changes r to $-t = s$ (from (22)). Again, by antisymmetry of $\delta\psi_0$, $r = t = -s$. As X, Y, Z are arbitrary, one gets $\psi_0 \in \mathfrak{c} \otimes (S^2 \mathfrak{g}^*)^{\mathfrak{g}}$. Now $F(\psi_0) = -r = -\delta\psi_0 = d\psi_1 \in B^3(\mathfrak{g}, \mathfrak{g})$. Hence

$$\psi_0 \in ZL_0^2(\mathfrak{g}, \mathfrak{g}) \Leftrightarrow F(\psi_0) = 0 \Leftrightarrow \psi_1 \in Z^2(\mathfrak{g}, \mathfrak{g}) \Leftrightarrow \psi_0 \in \mathfrak{c} \otimes \ker \mathcal{I}.$$

Consider now the linear map $\Phi : ZL^2(\mathfrak{g}, \mathfrak{g}) \rightarrow F^{-1}(B^3(\mathfrak{g}, \mathfrak{g})) / \ker F$ defined by $\psi \mapsto [\psi_0] \pmod{\ker F}$. Φ is onto: for any $[\varphi_0] \in F^{-1}(B^3(\mathfrak{g}, \mathfrak{g})) / \ker F$, $\varphi_0 \in \mathfrak{c} \otimes (S^2 \mathfrak{g}^*)^{\mathfrak{g}}$, one has $F(\varphi_0) \in B^3(\mathfrak{g}, \mathfrak{g})$, hence $F(\varphi_0) = d\varphi_1$, $\varphi_1 \in C^2(\mathfrak{g}, \mathfrak{g})$, and then $\varphi = \varphi_0 + \varphi_1$ is a Leibniz cocycle such that $\Phi(\varphi) = [\varphi_0]$. Now $\ker \Phi = Z^2(\mathfrak{g}, \mathfrak{g}) \oplus ZL_0^2(\mathfrak{g}, \mathfrak{g})$, since condition $[\psi_0] = [0]$ reads $\psi_0 \in \ker F$ which is equivalent to $\psi \in Z^2(\mathfrak{g}, \mathfrak{g}) \oplus ZL_0^2(\mathfrak{g}, \mathfrak{g})$. Hence Φ yields an isomorphism $ZL^2(\mathfrak{g}, \mathfrak{g}) / (Z^2(\mathfrak{g}, \mathfrak{g}) \oplus ZL_0^2(\mathfrak{g}, \mathfrak{g})) \cong F^{-1}(B^3(\mathfrak{g}, \mathfrak{g})) / \ker F$. The latter is isomorphic to $\text{Im } F \cap B^3(\mathfrak{g}, \mathfrak{g}) \cong (\mathfrak{c} \otimes \text{Im } \mathcal{I}) \cap B^3(\mathfrak{g}, \mathfrak{g})$.

(ii) Results from the invariance of $\psi_0 \in ZL_0^2(\mathfrak{g}, \mathfrak{g})$.

(iii) Results immediately from (i), (ii) since $BL^2(\mathfrak{g}, \mathfrak{g}) = B^2(\mathfrak{g}, \mathfrak{g})$ as the Leibniz differential on $CL^1(\mathfrak{g}, \mathfrak{g}) = \mathfrak{g}^* \otimes \mathfrak{g} = C^1(\mathfrak{g}, \mathfrak{g})$ coincides with the usual one.

(iv)-(vi) Readily similar. $\square$

Remark 4. Since $\ker \mathcal{I} \oplus (\text{Im } \mathcal{I} \cap B^3(\mathfrak{g}, \mathbb{C})) \cong \ker h$ where h denotes $\mathcal{I}$ composed with the projection of $Z^3(\mathfrak{g}, \mathbb{C})$ onto $H^3(\mathfrak{g}, \mathbb{C})$, the result (vi) is the same as in [10].

Remark 5. Any supplementary subspace to $Z^2(\mathfrak{g}, \mathbb{C}) \oplus ZL_0^2(\mathfrak{g}, \mathbb{C})$ in $ZL^2(\mathfrak{g}, \mathbb{C})$ consists of *coupled* Leibniz 2-cocycles, i.e. the nonzero elements have the property that their symmetric and antisymmetric parts are not cocycles. To get such a supplementary subspace, pick any supplementary subspace W to $\ker \mathcal{I}$ in $(S^2 \mathfrak{g}^*)^{\mathfrak{g}}$ and take $\mathcal{C} = \{B + \omega ; B \in W \cap \mathcal{I}^{-1}(B^3(\mathfrak{g}, \mathbb{C})), I_B = d\omega\}$.

Definition 2. $\mathfrak{g}$ is said to be *adjoint (resp. trivial) ZL^2 -uncoupling* if

$$(\mathfrak{c} \otimes \text{Im } \mathcal{I}) \cap B^3(\mathfrak{g}, \mathfrak{g}) = \{0\} \text{ (resp. } \text{Im } \mathcal{I} \cap B^3(\mathfrak{g}, \mathbb{C}) = \{0\}).$$

Adjoint ZL^2 -uncoupling implies trivial ZL^2 -uncoupling, since $\mathfrak{c} \otimes (\text{Im } \mathcal{I} \cap B^3(\mathfrak{g}, \mathbb{C})) \subset (\mathfrak{c} \otimes \text{Im } \mathcal{I}) \cap B^3(\mathfrak{g}, \mathfrak{g})$. The reciprocal holds obviously true for $\mathcal{I}$ -exact or zero-center Lie algebras. However we do not know if it holds true in general (e.g. we do not know of a nilpotent Lie algebra which is not $\mathcal{I}$ -exact). The class of adjoint ZL^2 -uncoupling Lie algebras is rather extensive since it contains all zero-center Lie algebras and all $\mathcal{I}$ -null Lie algebras.

Corollary 6. (i) $HL^2(\mathfrak{g}, \mathfrak{g}) \cong H^2(\mathfrak{g}, \mathfrak{g}) \oplus (\mathfrak{c} \otimes \ker \mathcal{I})$ if and only if $\mathfrak{g}$ is adjoint ZL^2 -uncoupling.

(ii) $HL^2(\mathfrak{g}, \mathbb{C}) \cong H^2(\mathfrak{g}, \mathbb{C}) \oplus \ker \mathcal{I}$ if and only if $\mathfrak{g}$ is trivial ZL^2 -uncoupling.

Example 6. $\mathfrak{g} = \mathfrak{g}_{5,4}$.

Trivial Leibniz cohomology. $B^2(\mathfrak{g}, \mathbb{C}) = \langle d\omega^3 = -\omega^{1,2}, d\omega^4 = -\omega^{1,3}, d\omega^5 = -\omega^{2,3} \rangle$, $\dim Z^2(\mathfrak{g}, \mathbb{C}) = 6$, $\dim H^2(\mathfrak{g}, \mathbb{C}) = 3$, $Z^2(\mathfrak{g}, \mathbb{C}) = \langle \omega^{1,4}, \omega^{2,5}, \omega^{1,5} + \omega^{2,4} \rangle \oplus B^2(\mathfrak{g}, \mathbb{C})$, $\dim ZL_0^2(\mathfrak{g}, \mathbb{C}) = 3$, $ZL_0^2(\mathfrak{g}, \mathbb{C}) (\cong \ker \mathcal{I}) = \langle \omega^1 \otimes \omega^1, \omega^1 \odot \omega^2, \omega^2 \otimes \omega^2 \rangle$, $\dim ZL^2(\mathfrak{g}, \mathbb{C}) = 10$, $\dim HL^2(\mathfrak{g}, \mathbb{C}) = 7$, and

$$\begin{aligned} ZL^2(\mathfrak{g}, \mathbb{C}) &= Z^2(\mathfrak{g}, \mathbb{C}) \oplus ZL_0^2(\mathfrak{g}, \mathbb{C}) \oplus \mathbb{C}g_1, \\ HL^2(\mathfrak{g}, \mathbb{C}) &= H^2(\mathfrak{g}, \mathbb{C}) \oplus ZL_0^2(\mathfrak{g}, \mathbb{C}) \oplus \mathbb{C}g_1 \end{aligned}$$

with $g_1 = B + \omega^{1,5}$ and $B = \omega^1 \odot \omega^5 - \omega^2 \odot \omega^4 + \omega^3 \otimes \omega^3$, (we already know that $\text{Im } \mathcal{I} = \mathbb{C}I_B = \mathbb{C}d\omega^{1,5}$ and $\text{Im } \mathcal{I} \cap B^3(\mathfrak{g}, \mathbb{C}) = \text{Im } \mathcal{I}$ is one-dimensional). $\mathfrak{g}_{5,4}$ is

not trivial ZL^2 -uncoupling (hence not adjoint ZL^2 -uncoupling either), and g_1 is a coupled Leibniz 2-cocycle.

Adjoint Leibniz cohomology. $\dim Z^2(\mathfrak{g}, \mathfrak{g}) = 24$; $ZL_0^2(\mathfrak{g}, \mathfrak{g}) = \mathfrak{c} \otimes \ker \mathcal{I}$ has dimension 6, $\dim ZL^2(\mathfrak{g}, \mathfrak{g}) = 32$,

$$\begin{aligned} ZL^2(\mathfrak{g}, \mathfrak{g}) &= Z^2(\mathfrak{g}, \mathfrak{g}) \oplus ZL_0^2(\mathfrak{g}, \mathfrak{g}) \oplus \mathbb{C}G_1 \oplus \mathbb{C}G_2, \\ HL^2(\mathfrak{g}, \mathfrak{g}) &= H^2(\mathfrak{g}, \mathfrak{g}) \oplus ZL_0^2(\mathfrak{g}, \mathfrak{g}) \oplus \mathbb{C}G_1 \oplus \mathbb{C}G_2, \end{aligned}$$

where G_1, G_2 are the following Leibniz 2-cocycles, each of which is coupled:

$$\begin{aligned} G_1 &= x_5 \otimes (B + \omega^{1,5}) \\ G_2 &= x_4 \otimes (B + \omega^{1,5}). \end{aligned}$$

Corollary 7. *For any Lie algebra $\mathfrak{g}$ with trivial center $\mathfrak{c} = \{0\}$, $HL^2(\mathfrak{g}, \mathfrak{g}) = H^2(\mathfrak{g}, \mathfrak{g})$. In particular any rigid Lie algebra with trivial center is Leibniz rigid.*

Corollary 8. *For any reductive algebra Lie $\mathfrak{g}$ with center $\mathfrak{c}$, $HL^2(\mathfrak{g}, \mathfrak{g}) = H^2(\mathfrak{g}, \mathfrak{g}) \oplus (\mathfrak{c} \otimes S^2\mathfrak{c}^*)$, and $\dim H^2(\mathfrak{g}, \mathfrak{g}) = \frac{c^2(c-1)}{2}$ with $c = \dim \mathfrak{c}$.*

Proof. $\mathfrak{g} = \mathfrak{s} \oplus \mathfrak{c}$ with $\mathfrak{s} = \mathcal{C}^2\mathfrak{g}$ semisimple. We first prove that $\mathfrak{g}$ is adjoint ZL^2 -uncoupling. $\mathfrak{c} \otimes (S^2\mathfrak{g}^*)^{\mathfrak{g}} = \mathfrak{c} \otimes (S^2\mathfrak{s}^*)^{\mathfrak{s}} \oplus (\mathfrak{c} \otimes S^2\mathfrak{c}^*) = c (S^2\mathfrak{s}^*)^{\mathfrak{s}} \oplus c (S^2\mathfrak{c}^*)$. Suppose first $\mathfrak{s}$ simple. Then any bilinear symmetric invariant form on $\mathfrak{s}$ is some multiple of the Killing form K . Hence $\mathfrak{c} \otimes (S^2\mathfrak{g}^*)^{\mathfrak{g}} = c(\mathbb{C}K) \oplus c(S^2\mathfrak{c}^*)$. For any $\psi_0 \in \mathfrak{c} \otimes (S^2\mathfrak{g}^*)^{\mathfrak{g}}$, $F(\psi_0)$ is then some linear combination of copies of I_K . As is well-known, I_K is no coboundary. Hence if we suppose that $F(\psi_0)$ is a coboundary, necessarily $F(\psi_0) = 0$. $\mathfrak{g}$ is adjoint ZL^2 -uncoupling when $\mathfrak{s}$ is simple. Now, if $\mathfrak{s}$ is not simple, $\mathfrak{s}$ can be decomposed as a direct sum $\mathfrak{s}_1 \oplus \dots \oplus \mathfrak{s}_m$ of simple ideals of $\mathfrak{s}$. Then $(S^2\mathfrak{s}^*)^{\mathfrak{s}} = \bigoplus_{i=1}^m (S^2\mathfrak{s}_i^*)^{\mathfrak{s}_i} = \bigoplus_{i=1}^m \mathbb{C}K_i$ (K_i Killing form of $\mathfrak{s}_i$). The same reasoning then applies and shows that $\mathfrak{g}$ is adjoint ZL^2 -uncoupling. From (ii) in the Theorem, $ZL_0^2(\mathfrak{g}, \mathfrak{g}) = \mathfrak{c} \otimes S^2\mathfrak{c}^*$. Now, $\mathfrak{g} = \mathfrak{s} \oplus \mathfrak{c}$ with $\mathfrak{s} = \mathcal{C}^2\mathfrak{g}$ semisimple. $\mathfrak{s}$ can be decomposed as a direct sum $\mathfrak{s}_1 \oplus \dots \oplus \mathfrak{s}_m$ of ideals of $\mathfrak{s}$ hence of $\mathfrak{g}$. Then $H^2(\mathfrak{g}, \mathfrak{g}) = \bigoplus_{i=1}^m H^2(\mathfrak{g}, \mathfrak{s}_i) \oplus H^2(\mathfrak{g}, \mathfrak{c})$. As $\mathfrak{s}_i$ is a nontrivial $\mathfrak{g}$ -module, $H^2(\mathfrak{g}, \mathfrak{s}_i) = \{0\}$ ([5], Prop. 11.4, page 154). Hence $H^2(\mathfrak{g}, \mathfrak{g}) = H^2(\mathfrak{g}, \mathfrak{c}) = c H^2(\mathfrak{g}, \mathbb{C})$. By the Künneth formula and Whitehead's lemmas, $H^2(\mathfrak{g}, \mathbb{C}) = (H^2(\mathfrak{s}, \mathbb{C}) \otimes H^0(\mathfrak{c}, \mathbb{C})) \oplus (H^1(\mathfrak{s}, \mathbb{C}) \otimes H^1(\mathfrak{c}, \mathbb{C})) \oplus (H^0(\mathfrak{s}, \mathbb{C}) \otimes H^2(\mathfrak{c}, \mathbb{C})) = H^0(\mathfrak{s}, \mathbb{C}) \otimes H^2(\mathfrak{c}, \mathbb{C}) = \mathbb{C} \otimes H^2(\mathfrak{c}, \mathbb{C})$. Hence $\dim H^2(\mathfrak{g}, \mathfrak{g}) = \frac{c^2(c-1)}{2}$. $\square$

Example 7. For $\mathfrak{g} = \mathfrak{gl}(n)$, $HL^2(\mathfrak{g}, \mathfrak{g}) = ZL_0^2(\mathfrak{g}, \mathfrak{g}) = \mathbb{C} \left(x_{n^2} \oplus (\omega^{n^2} \circ \omega^{n^2}) \right)$, where $(x_i)_{1 \leq i \leq n^2}$ is a basis of $\mathfrak{g}$ such that $(x_i)_{1 \leq i \leq n^2-1}$ is a basis of $\mathfrak{sl}(n)$ and x_{n^2} is the identity matrix, and $(\omega_i)_{1 \leq i \leq n^2}$ the dual basis to $(x_i)_{1 \leq i \leq n^2}$. Hence there is a unique Leibniz deformation of $\mathfrak{gl}(n)$.

Corollary 9. *Let $\mathcal{H}_N$ be the $(2N+1)$ -dimensional complex Heisenberg Lie algebra ($N \geq 1$) with basis $(x_i)_{1 \leq i \leq 2N+1}$ and commutation relations $[x_i, x_{N+i}] = x_{2N+1}$ ($1 \leq i \leq N$).*

(i) $ZL_0^2(\mathcal{H}_N, \mathcal{H}_N)$ has basis $(x_{2N+1} \otimes (\omega^i \circ \omega^j))_{1 \leq i \leq j \leq 2N}$ with $(\omega_i)_{1 \leq i \leq 2N+1}$ the dual basis to $(x_i)_{1 \leq i \leq 2N+1}$.

(ii)

$$\dim ZL_0^2(\mathcal{H}_N, \mathcal{H}_N) = \dim B^2(\mathcal{H}_N, \mathcal{H}_N) = N(2N+1);$$

$$\dim HL^2(\mathcal{H}_N, \mathcal{H}_N) = \dim Z^2(\mathcal{H}_N, \mathcal{H}_N) = \begin{cases} \frac{N}{3}(8N^2 + 6N + 1) & \text{if } N \geq 2 \\ 8 & \text{if } N = 1. \end{cases}$$

Proof. (i) Follows from $\ker \mathcal{I} = S^2(\mathfrak{g}/\mathcal{C}^2\mathfrak{g})^*$.

(ii) First $\mathcal{H}_N$ is adjoint ZL^2 -uncoupling since it is $\mathcal{I}$ -null. The result then follows from the fact that ([12]) $\dim B^2(\mathcal{H}_N, \mathcal{H}_N) = N(2N+1)$ and for $N \geq 2$, $\dim H^2(\mathcal{H}_N, \mathcal{H}_N) = \frac{2N}{3}(4N^2 - 1)$. $\square$

Example 8. The case $N = 1$ has been studied in [4]. In that case, $\dim ZL_0^2(\mathcal{H}_1, \mathcal{H}_1) = 3$ and the 3 Leibniz deformations are nilpotent, in contradistinction with the 5 Lie deformations.

References

- [1] G. Favre, L. J. Santharoubane, Nilpotent Lie algebras of classical simple type *J. Algebra*, **202**, #2, 1998, 589-910.
- [2] D. Fernandez-Ternero, Nilpotent Lie algebras of maximal rank and of Kac-Moody type $D_4^{(3)}$, *J. of Lie Theory*, **15**, #1, 2005, 249-260.
- [3] D. Fernandez-Ternero, J. Nunez-Valdes, Nilpotent Lie algebras of maximal rank and of Kac-Moody type $F_4^{(1)}$, *Comm. Algebra*, **29**, #4, 2001, 1151-1570.
- [4] A. Fialowski, A. Mandal, Leibniz algebra deformations of a Lie algebra *preprint ArXiv 0802.1263*, 2008.
- [5] A. Guichardet, *Cohomologie des groupes topologiques et des algèbres de Lie*, Cedric/Fernand Nathan, Paris, 1980.
- [6] W. Fulton, J. Harris *Representation Theory*, Grad. Texts in Math. #129, Springer-Verlag, New York, 1991.
- [7] S. Helgason, *Differential Geometry, Lie groups, and symmetric spaces*, Academic Press, New York, 1978.
- [8] V. Kac, *Infinite dimensional Lie algebras*, 3^d edition, Cambridge University Press, 1990.
- [9] J.L. Koszul, Homologie et cohomologie des algèbres de Lie, *Bull. Soc. Math. France*, **78**, 1950, 67-127.

- [10] Naihong Hu, Yufeng Pei, Dong Liu A cohomological characterization of Leibniz central extensions of Lie algebras, *Proc. Amer. Math. Soc.*, **136**, 2008, 437-477.
- [11] J.L. Loday, Une version non commutative des algèbres de Lie: les algèbres de Leibniz, *Ens. Math.*, **39**, 1993, 269-293.
- [12] L. Magnin, Cohomologie adjointe de algèbres de Heisenberg, *Comm. Algebra*, **21**, 1993, 2101-2129.
- [13] L. Magnin, Adjoint and trivial cohomologies of nilpotent complex Lie algebras of dimension ≤ 7 , *Int. J. Math. math. Sci.*, volume 2008, Article ID 805305, 12 pages.
- [14] L. Magnin, Determination of 7-dimensional indecomposable nilpotent complex Lie algebras by adjoining a derivation to 6-dimensional Lie algebras, *Algebras and Representation Theory*, to appear, <http://www.u-bourgogne.fr/monge/l.magnin> or <http://math.u-bourgogne.fr/IMB/magnin/public.html/index.html>
- [15] L. Magnin, *Adjoint and trivial cohomology tables for indecomposable nilpotent Lie algebras of dimension ≤ 7 over $\mathbb{C}$* , online book, 2d Corrected Edition 2007, (*Postscript, .ps file*) (810 pages + vi), accessible at <http://www.u-bourgogne.fr/monge/l.magnin> or <http://math.u-bourgogne.fr/IMB/magnin/public.html/index.html>
- [16] <http://www.u-bourgogne.fr/monge/l.magnin/CL/CLindex.html> or <http://math.u-bourgogne.fr/IMB/magnin/public.html/CLindex.html>
- [17] A. Medina, Groupes de Lie munis de pseudo-métriques de Riemann bi-invariantes, *Séminaire de Géométrie différentielle, exposé #6*, Montpellier, 1982.
- [18] G. Ovando, Complex, symplectic and Kähler structures on 4 dimensional Lie groups, *preprint ArXiv DG/0309146*, 2003.
- [19] T. Pirashvili, On Leibniz homology, *Ann. Instit. Fourier*, **44**, 1994, 401-411.
- [20] A. Medina, P. Revoy, Algèbres de Lie et produit scalaire invariant, *Ann. Scient. Ec. Norm. Sup*, **18**, 1985, 553-561.
- [21] L. J. Santharoubane, Kac-Moody Lie algebras and the universal element for the category of nilpotent Lie algebras, *Math. Ann.*, **263**, 1983, 365-370.
- [22] L. J. Santharoubane, Nilpotent Lie algebras of Kac-Moody affine type, *J. Algebra*, **302**, #2, 2006, 553-585.
- [23] F. Wagemann, Habilitation, 2007, <http://www.math.sciences.univ-nantes.fr/~wagemann/hdr.pdf>
- [24] Wan Zhe-Xian, *Introduction to Kac-Moody algebras*, World Scientific, Singapore, 1991.

Table 2: Kac-Moody types for indecomposable nilpotent Lie algebras of dimension ≤ 7 . Notations for indefinite hyperbolic are those of [24].

algebra	GCM	Finite	Affine	Indefinite Hyperbolic	Indefinite Not Hyperbolic
$\mathfrak{g}_3$	$\begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$	A_1			
$\mathfrak{g}_4$	$\begin{pmatrix} 2 & -2 \\ -1 & 2 \end{pmatrix}$	C_2			
$\mathfrak{g}_{5,1}$	$\begin{pmatrix} 2 & 0 & -1 & 0 \\ 0 & 2 & 0 & -1 \\ -1 & 0 & 2 & 0 \\ 0 & -1 & 0 & 2 \end{pmatrix}$	$A_2 \times A_2$			
$\mathfrak{g}_{5,2}$	$\begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}$	A_3			
$\mathfrak{g}_{5,3}$	$\begin{pmatrix} 2 & -2 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}$	B_3			
$\mathfrak{g}_{5,4}$	$\begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$		$A_1^{(1)}$		
$\mathfrak{g}_{5,5}$	$\begin{pmatrix} 2 & -3 \\ -1 & 2 \end{pmatrix}$	G_2			
$\mathfrak{g}_{5,6}$	$\begin{pmatrix} 2 & -3 \\ -2 & 2 \end{pmatrix}$			$(3, 2)$	
$\mathfrak{g}_{6,1}$	$\begin{pmatrix} 2 & -1 & 0 & -1 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & 0 \\ -1 & 0 & 0 & 2 \end{pmatrix}$	D_4			
$\mathfrak{g}_{6,2}$	$\begin{pmatrix} 2 & -2 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & 0 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{pmatrix}$	$B_2 \times A_2$			
$\mathfrak{g}_{6,3}$	$\begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$		$A_2^{(1)}$		
$\mathfrak{g}_{6,4}$	$\begin{pmatrix} 2 & -1 & -1 \\ -2 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}$	B_3			
$\mathfrak{g}_{6,5}$	$\begin{pmatrix} 2 & -2 & -1 \\ -2 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$			$H_2^{(3)}$	
$\mathfrak{g}_{6,6}$	$\begin{pmatrix} 2 & -1 & 0 \\ -2 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}$	C_3			
$\mathfrak{g}_{6,7}$	$\begin{pmatrix} 2 & -2 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$			$H_1^{(3)}$	
$\mathfrak{g}_{6,8}$	$\begin{pmatrix} 2 & -2 & 0 \\ -2 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}$			$H_{96}^{(3)}$	
$\mathfrak{g}_{6,9}$	$\begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}$	A_3			
$\mathfrak{g}_{6,10}$	$\begin{pmatrix} 2 & -2 & -1 \\ -1 & 2 & 0 \\ -2 & 0 & 2 \end{pmatrix}$		$A_4^{(2)}$		
$\mathfrak{g}_{6,11}$	$\begin{pmatrix} 2 & -3 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}$		$G_2^{(1)}$		
$\mathfrak{g}_{6,12}$	$\begin{pmatrix} 2 & -3 & -2 \\ -2 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$			$H_{10}^{(3)}$	
$\mathfrak{g}_{6,13}$	$\begin{pmatrix} 2 & -1 & 0 \\ -3 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}$		$G_2^{(1)}$		
$\mathfrak{g}_{6,14}$	$\begin{pmatrix} 2 & -3 \\ -2 & 2 \end{pmatrix}$			$(3, 2)$	
$\mathfrak{g}_{6,15}$	$\begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$		$A_1^{(1)}$		
$\mathfrak{g}_{6,16}$	$\begin{pmatrix} 2 & -4 \\ -1 & 2 \end{pmatrix}$		$A_2^{(2)}$		
$\mathfrak{g}_{6,17}$	$\begin{pmatrix} 2 & -4 \\ -2 & 2 \end{pmatrix}$			$(4, 2)$	

Table 2: continued

algebra	GCM	Finite	Affine	Indefinite Hyperbolic	Indefinite Not Hyper- bolic
$\mathfrak{g}_{6,18}$	$\begin{pmatrix} 2 & -3 \\ -1 & 2 \end{pmatrix}$	G_2	$A_2^{(2)}$		
$\mathfrak{g}_{6,19}$	$\begin{pmatrix} 2 & -4 \\ -1 & 2 \end{pmatrix}$				
$\mathfrak{g}_{6,20}$	$\begin{pmatrix} 2 & -3 \\ -3 & 2 \end{pmatrix}$			(3, 3)	
$\mathfrak{g}_{7,0.1}$	$\begin{pmatrix} 2 & -5 \\ -5 & 2 \end{pmatrix}$			(5, 5)	
$\mathfrak{g}_{7,0.2}$	idem			idem	
$\mathfrak{g}_{7,0.3}$	idem			idem	
$\mathfrak{g}_{7,0.4(\lambda)}$	$\begin{pmatrix} 2 & -4 \\ -4 & 2 \end{pmatrix}$			(4, 4)	
$\mathfrak{g}_{7,0.5}$	idem			idem	
$\mathfrak{g}_{7,0.6}$	$\begin{pmatrix} 2 & -3 \\ -3 & 2 \end{pmatrix}$			(3, 3)	
$\mathfrak{g}_{7,0.7}$	idem			idem	
$\mathfrak{g}_{7,0.8}$	$\begin{pmatrix} 2 & -3 & -3 \\ -3 & 2 & -3 \\ -3 & -3 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,1.01(i)}$	$\begin{pmatrix} 2 & 0 & -4 \\ 0 & 2 & -4 \\ -1 & -1 & 2 \end{pmatrix}$			$H_{123}^{(3)}$	
$\mathfrak{g}_{7,1.01(ii)}$	idem			idem	
$\mathfrak{g}_{7,1.02}$	$\begin{pmatrix} 2 & -2 \\ -3 & 2 \end{pmatrix}$			(3, 2)	
$\mathfrak{g}_{7,1.03}$	$\begin{pmatrix} 2 & -3 \\ -2 & 2 \end{pmatrix}$			(3, 2)	
$\mathfrak{g}_{7,1.1(i_\lambda)}$ $\lambda \neq 0$	$\begin{pmatrix} 2 & -5 \\ -3 & 2 \end{pmatrix}$			(5, 3)	
$\mathfrak{g}_{7,1.1(i_\lambda)}$ $\lambda = 0$	$\begin{pmatrix} 2 & -5 \\ -2 & 2 \end{pmatrix}$			(5, 2)	
$\mathfrak{g}_{7,1.1(ii)}$	$\begin{pmatrix} 2 & -5 \\ -1 & 2 \end{pmatrix}$			(5, 1)	
$\mathfrak{g}_{7,1.1(iii)}$	$\begin{pmatrix} 2 & -4 \\ -3 & 2 \end{pmatrix}$			(4, 3)	
$\mathfrak{g}_{7,1.1(iv)}$	$\begin{pmatrix} 2 & -2 \\ -3 & 2 \end{pmatrix}$			(4, 2)	
$\mathfrak{g}_{7,1.1(v)}$	$\begin{pmatrix} 2 & 0 & -4 \\ 0 & 2 & -2 \\ -2 & -1 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,1.1(vi)}$	$\begin{pmatrix} 2 & -4 & -1 \\ -3 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,1.2(i_\lambda)}$	$\begin{pmatrix} 2 & -3 & -2 \\ -3 & 2 & -2 \\ -1 & -1 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,1.2(ii)}$	$\begin{pmatrix} 2 & -3 & -2 \\ -3 & 2 & -2 \\ -1 & -1 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,1.2(iii)}$	$\begin{pmatrix} 2 & -3 \\ -2 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,1.2(iv)}$	$\begin{pmatrix} 2 & -3 & -2 \\ -3 & 2 & -2 \\ -1 & -1 & 2 \end{pmatrix}$				✓

Table 2: continued

algebra	GCM	Finite	Affine	Indefinite Hyperbolic	Indefinite Not Hyper- bolic
$\mathfrak{g}_{7,1.3(i\lambda)}$	$\begin{pmatrix} 2 & -3 & -3 \\ -2 & 2 & -1 \\ -2 & -1 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,1.3(ii)}$	idem				idem
$\mathfrak{g}_{7,1.3(iii)}$	$\begin{pmatrix} 2 & -3 & -3 \\ -2 & 2 & 0 \\ -2 & 0 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,1.3(iv)}$	$\begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -1 \\ -2 & -1 & 2 \end{pmatrix}$			$H_{18}^{(3)}$	
$\mathfrak{g}_{7,1.3(v)}$	$\begin{pmatrix} 2 & -3 & -3 & -2 \\ -2 & 2 & -1 & -1 \\ -2 & -1 & 2 & -1 \\ -2 & -1 & -1 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,1.4}$	$\begin{pmatrix} 2 & -5 \\ -2 & 2 \end{pmatrix}$			$(5, 2)$	
$\mathfrak{g}_{7,1.5}$	$\begin{pmatrix} 2 & -4 \\ -2 & 2 \end{pmatrix}$			$(4, 2)$	
$\mathfrak{g}_{7,1.6}$	$\begin{pmatrix} 2 & -5 \\ -2 & 2 \end{pmatrix}$			$(5, 2)$	
$\mathfrak{g}_{7,1.7}$	$\begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$			$H_8^{(3)}$	
$\mathfrak{g}_{7,1.8}$	$\begin{pmatrix} 2 & -3 & 0 \\ -2 & 2 & -1 \\ 0 & -2 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,1.9}$	$\begin{pmatrix} 2 & -3 & -1 \\ -2 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,1.10}$	$\begin{pmatrix} 2 & -4 \\ -3 & 2 \end{pmatrix}$			$(4, 3)$	
$\mathfrak{g}_{7,1.11}$	$\begin{pmatrix} 2 & -4 & -3 \\ -2 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,1.12}$	$\begin{pmatrix} 2 & -4 & -2 \\ -2 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,1.13}$	$\begin{pmatrix} 2 & -4 \\ -2 & 2 \end{pmatrix}$			$(4, 2)$	
$\mathfrak{g}_{7,1.14}$	$\begin{pmatrix} 2 & -3 \\ -3 & 2 \end{pmatrix}$			$(3, 3)$	
$\mathfrak{g}_{7,1.15}$	$\begin{pmatrix} 2 & -4 & -3 \\ -2 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,1.16}$	$\begin{pmatrix} 2 & -3 & -2 \\ -2 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,1.17}$	$\begin{pmatrix} 2 & -4 \\ -4 & 2 \end{pmatrix}$			$(4, 4)$	
$\mathfrak{g}_{7,1.18}$	$\begin{pmatrix} 2 & -3 & -1 \\ -2 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,1.19}$	$\begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix}$			$H_{71}^{(3)}$	
$\mathfrak{g}_{7,1.20}$	$\begin{pmatrix} 2 & -1 & 0 \\ -3 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}$		$D_4^{(3)}$		
$\mathfrak{g}_{7,1.21}$	$\begin{pmatrix} 2 & -3 & -2 \\ -3 & 2 & -2 \\ -1 & -1 & 2 \end{pmatrix}$				✓

Table 2: continued

algebra	GCM	Finite	Affine	Indefinite Hyperbolic	Indefinite Not Hyper- bolic
$\mathfrak{g}_{7,2.1(i\lambda)}$	$\begin{pmatrix} 2 & -3 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$			$H_3^{(3)}$	
$\mathfrak{g}_{7,2.1(ii)}$	$\begin{pmatrix} 2 & -3 & -1 \\ -1 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}$		$D_4^{(3)}$		
$\mathfrak{g}_{7,2.1(iii)}$	$\begin{pmatrix} 2 & -3 & 0 & 0 \\ -1 & 2 & -1 & -1 \\ 0 & -1 & 2 & 0 \\ 0 & -1 & 0 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,2.1(iv)}$	$\begin{pmatrix} 2 & 0 & -1 & -2 \\ 0 & 2 & -1 & 0 \\ -1 & -1 & 2 & -1 \\ -2 & 0 & -1 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,2.1(v)}$	$\begin{pmatrix} 2 & -2 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$			$H_1^{(3)}$	
$\mathfrak{g}_{7,2.2}$	$\begin{pmatrix} 2 & -1 & -1 \\ -2 & 2 & -1 \\ -2 & -1 & 2 \end{pmatrix}$			$H_7^{(3)}$	
$\mathfrak{g}_{7,2.3}$	$\begin{pmatrix} 2 & -5 \\ -2 & 2 \end{pmatrix}$			$(5, 1)$	
$\mathfrak{g}_{7,2.4}$	$\begin{pmatrix} 2 & -4 \\ -1 & 2 \end{pmatrix}$		$A_2^{(2)}$		
$\mathfrak{g}_{7,2.5}$	$\begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$		$A_1^{(1)}$		
$\mathfrak{g}_{7,2.6}$	$\begin{pmatrix} 2 & -3 \\ -2 & 2 \end{pmatrix}$			$(3, 2)$	
$\mathfrak{g}_{7,2.7}$	$\begin{pmatrix} 2 & -4 \\ -2 & 2 \end{pmatrix}$			$(4, 2)$	
$\mathfrak{g}_{7,2.8}$	$\begin{pmatrix} 2 & -3 \\ -2 & 2 \end{pmatrix}$			$(3, 2)$	
$\mathfrak{g}_{7,2.9}$	$\begin{pmatrix} 2 & -3 \\ -3 & 2 \end{pmatrix}$			$(3, 3)$	
$\mathfrak{g}_{7,2.10}$	$\begin{pmatrix} 2 & -3 & -1 \\ -1 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}$		$D_4^{(3)}$		
$\mathfrak{g}_{7,2.11}$	$\begin{pmatrix} 2 & -3 & -2 \\ -2 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,2.12}$	$\begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & 0 \\ -2 & 0 & 2 \end{pmatrix}$			$H_{109}^{(3)}$	
$\mathfrak{g}_{7,2.13}$	$\begin{pmatrix} 2 & -3 & 0 \\ -1 & 2 & -2 \\ 0 & -1 & 2 \end{pmatrix}$			$H_{100}^{(3)}$	
$\mathfrak{g}_{7,2.14}$	$\begin{pmatrix} 2 & -4 & 0 \\ -1 & 2 & -1 \\ 0 & -2 & 2 \end{pmatrix}$			$H_{107}^{(3)}$	
$\mathfrak{g}_{7,2.15}$	$\begin{pmatrix} 2 & -4 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}$			$H_{97}^{(3)}$	
$\mathfrak{g}_{7,2.16}$	idem			idem	
$\mathfrak{g}_{7,2.17}$	$\begin{pmatrix} 2 & -3 & 0 \\ -2 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,2.18}$	idem				idem
$\mathfrak{g}_{7,2.19}$	$\begin{pmatrix} 2 & -3 & -1 \\ -2 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,2.20}$	$\begin{pmatrix} 2 & -1 & -3 \\ -1 & 2 & 0 \\ -2 & 0 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,2.21}$	$\begin{pmatrix} 2 & -3 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$			$H_1^{(3)}$	
$\mathfrak{g}_{7,2.22}$	$\begin{pmatrix} 2 & 0 & -3 \\ 0 & 2 & -1 \\ -2 & -1 & 2 \end{pmatrix}$				✓
$\mathfrak{g}_{7,2.23}$	$\begin{pmatrix} 2 & 0 & 0 & -2 \\ 0 & 2 & -2 & 0 \\ 0 & -1 & 2 & -1 \\ -1 & 0 & -1 & 2 \end{pmatrix}$		$D_4^{(2)}$		

Table 2: continued

algebra	GCM	Finite	Affine	Indefinite Hyperbolic	Indefinite Not Hyper- bolic
$\mathfrak{g}_{7,2.24}$	$\begin{pmatrix} 2 & -3 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}$		$G_2^{(1)}$		
$\mathfrak{g}_{7,2.25}$	$\begin{pmatrix} 2 & -3 & 0 & -2 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ -1 & 0 & -1 & 2 \end{pmatrix}$				$\checkmark$
$\mathfrak{g}_{7,2.26}$	$\begin{pmatrix} 2 & -1 & -2 \\ -1 & 2 & -1 \\ -2 & -1 & 2 \end{pmatrix}$			$H_8^{(3)}$	
$\mathfrak{g}_{7,2.27}$	$\begin{pmatrix} 2 & -2 & -1 & -1 \\ -2 & 2 & 0 & -1 \\ -1 & 0 & 2 & 0 \\ -1 & -1 & 0 & 2 \end{pmatrix}$				$\checkmark$
$\mathfrak{g}_{7,2.28}$	$\begin{pmatrix} 2 & -1 & -2 & 0 \\ -2 & 2 & 0 & 0 \\ -1 & 0 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{pmatrix}$			$H_{40}^{(4)}$	
$\mathfrak{g}_{7,2.29}$	$\begin{pmatrix} 2 & -2 & 0 & 0 \\ -2 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{pmatrix}$				$\checkmark$
$\mathfrak{g}_{7,2.30}$	$\begin{pmatrix} 2 & -3 & 0 & 0 \\ -2 & 2 & 0 & 0 \\ 0 & 0 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{pmatrix}$			$(3, 2) \times A_1$	
$\mathfrak{g}_{7,2.31}$	$\begin{pmatrix} 2 & -3 & 0 \\ -1 & 2 & -2 \\ 0 & -1 & 2 \end{pmatrix}$			$H_{100}^{(3)}$	
$\mathfrak{g}_{7,2.32}$	$\begin{pmatrix} 2 & -3 & -1 \\ -1 & 2 & 0 \\ -2 & 0 & 2 \end{pmatrix}$			$H_{106}^{(3)}$	
$\mathfrak{g}_{7,2.33}$	$\begin{pmatrix} 2 & -3 & 0 \\ -1 & 2 & -1 \\ 0 & -2 & 2 \end{pmatrix}$			$H_{105}^{(3)}$	
$\mathfrak{g}_{7,2.34}$	$\begin{pmatrix} 2 & -2 & -1 \\ -2 & 2 & 0 \\ -2 & 0 & 2 \end{pmatrix}$			$H_{104}^{(3)}$	
$\mathfrak{g}_{7,2.35}$	$\begin{pmatrix} 2 & -1 & -2 \\ -2 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}$		$A_4^{(2)}$		
$\mathfrak{g}_{7,2.36}$	$\begin{pmatrix} 2 & 0 & -1 & -1 \\ 0 & 2 & -1 & -1 \\ -1 & -2 & 2 & 0 \\ -1 & -1 & 0 & 2 \end{pmatrix}$				$\checkmark$
$\mathfrak{g}_{7,2.37}$	$\begin{pmatrix} 2 & -2 & -1 \\ -2 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}$			$H_{96}^{(3)}$	
$\mathfrak{g}_{7,2.38}$	$\begin{pmatrix} 2 & -2 & 0 & -1 \\ -2 & 2 & -1 & 0 \\ 0 & -1 & 2 & 0 \\ -1 & 0 & 0 & 2 \end{pmatrix}$				$\checkmark$
$\mathfrak{g}_{7,2.39}$	$\begin{pmatrix} 2 & -2 & -2 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$			$H_5^{(3)}$	
$\mathfrak{g}_{7,2.40}$	$\begin{pmatrix} 2 & -2 & -1 \\ -2 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$			$H_2^{(3)}$	
$\mathfrak{g}_{7,2.41}$	$\begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}$			$H_{99}^{(3)}$	
$\mathfrak{g}_{7,2.42}$	$\begin{pmatrix} 2 & -1 & -2 \\ -2 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$			$H_6^{(3)}$	
$\mathfrak{g}_{7,2.43}$	$\begin{pmatrix} 2 & -2 & -2 \\ -1 & 2 & 0 \\ -2 & 0 & 2 \end{pmatrix}$			$H_{99}^{(3)}$	
$\mathfrak{g}_{7,2.44}$	$\begin{pmatrix} 2 & -1 & -2 \\ -2 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$			$H_6^{(3)}$	
$\mathfrak{g}_{7,2.45}$	$\begin{pmatrix} 2 & -2 & 0 & -1 \\ -1 & 2 & -1 & -1 \\ 0 & -1 & 2 & 0 \\ -1 & -1 & 0 & 2 \end{pmatrix}$				$\checkmark$

Table 2: continued

algebra	GCM	Finite	Affine	Indefinite Hyperbolic	Indefinite Not Hyper- bolic
$\mathfrak{g}_{7,3.1(i_\lambda)}$	$\begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$	D_4	$A_2^{(1)}$	$H_1^{(3)}$	
$\mathfrak{g}_{7,3.1(iii)}$	$\begin{pmatrix} 2 & -1 & -1 & -1 \\ -1 & 2 & 0 & 0 \\ -1 & 0 & 2 & 0 \\ -1 & 0 & 0 & 2 \end{pmatrix}$				
$\mathfrak{g}_{7,3.2}$	$\begin{pmatrix} 2 & -3 & -1 \\ -1 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}$		$D_4^{(3)}$		
$\mathfrak{g}_{7,3.3}$	$\begin{pmatrix} 2 & -3 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}$	B_3	$G_2^{(1)}$		
$\mathfrak{g}_{7,3.4}$	$\begin{pmatrix} 2 & -1 & -1 \\ -2 & 2 & 0 \\ -2 & 0 & 2 \end{pmatrix}$		$D_3^{(2)}$		
$\mathfrak{g}_{7,3.5}$	$\begin{pmatrix} 2 & -1 & -1 \\ -2 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}$				
$\mathfrak{g}_{7,3.6}$	$\begin{pmatrix} 2 & -1 & -2 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$	B_4			
$\mathfrak{g}_{7,3.7}$	$\begin{pmatrix} 2 & -2 & 0 & 0 \\ -1 & 2 & 0 & -1 \\ 0 & 0 & 2 & -1 \\ 0 & -1 & -1 & 2 \end{pmatrix}$				
$\mathfrak{g}_{7,3.8}$	$\begin{pmatrix} 2 & -2 & -1 & 0 \\ -1 & 2 & 0 & -1 \\ -1 & 0 & 2 & 0 \\ 0 & -1 & 0 & 2 \end{pmatrix}$				
$\mathfrak{g}_{7,3.9}$	$\begin{pmatrix} 2 & -2 & 0 & 0 \\ -1 & 2 & -1 & -1 \\ 0 & -1 & 2 & 0 \\ 0 & -1 & 0 & 2 \end{pmatrix}$	F_4	$B_4^{(1)}$		
$\mathfrak{g}_{7,3.10}$	$\begin{pmatrix} 2 & -1 & -1 & 0 \\ -1 & 2 & 0 & -2 \\ -1 & 0 & 2 & 0 \\ 0 & -1 & 0 & 2 \end{pmatrix}$				
$\mathfrak{g}_{7,3.11}$	$\begin{pmatrix} 2 & -2 & -1 & 0 \\ -1 & 2 & 0 & -1 \\ -1 & 0 & 2 & 0 \\ 0 & -1 & 0 & 2 \end{pmatrix}$				
$\mathfrak{g}_{7,3.12}$	$\begin{pmatrix} 2 & -1 & -1 & 0 \\ -1 & 2 & 0 & -1 \\ -1 & 0 & 2 & -1 \\ 0 & -1 & -1 & 2 \end{pmatrix}$	C_4	$A_3^{(1)}$		
$\mathfrak{g}_{7,3.13}$	$\begin{pmatrix} 2 & -2 & 0 & 0 \\ -2 & 2 & 0 & 0 \\ 0 & 0 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{pmatrix}$		$A_1^{(1)} \times A_1$		
$\mathfrak{g}_{7,3.14}$	$\begin{pmatrix} 2 & -1 & -2 & 0 \\ -1 & 2 & 0 & -1 \\ -1 & 0 & 2 & 0 \\ 0 & -1 & 0 & 2 \end{pmatrix}$				
$\mathfrak{g}_{7,3.15}$	$\begin{pmatrix} 2 & -1 & -1 & 0 \\ -2 & 2 & 0 & 0 \\ -1 & 0 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{pmatrix}$	B_4			
$\mathfrak{g}_{7,3.16}$	$\begin{pmatrix} 2 & -2 & 0 & 0 \\ -1 & 2 & 0 & 0 \\ 0 & 0 & 2 & -2 \\ 0 & 0 & -1 & 2 \end{pmatrix}$	$B_2 \times B_2$			
$\mathfrak{g}_{7,3.17}$	$\begin{pmatrix} 2 & -3 & 0 & 0 \\ -1 & 2 & 0 & 0 \\ 0 & 0 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{pmatrix}$	$G_2 \times A_2$			
$\mathfrak{g}_{7,3.18}$	$\begin{pmatrix} 2 & -2 & 0 & 0 & 0 \\ -1 & 2 & 0 & 0 & -1 \\ 0 & 0 & 2 & -1 & 0 \\ 0 & 0 & -1 & 2 & 0 \\ 0 & -1 & 0 & 0 & 2 \end{pmatrix}$	$B_3 \times A_2$			
$\mathfrak{g}_{7,3.19}$	$\begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & 0 & -1 & 0 \\ 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & -1 & 2 \end{pmatrix}$	A_5			

Table 2: continued

algebra	GCM	Finite	Affine	Indefinite Hyperbolic	Indefinite Not Hyper- bolic
$\mathfrak{g}_{7,3.20}$	$\begin{pmatrix} 2 & -2 & -2 \\ -1 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}$	C_3	$C_2^{(1)}$	$H_{96}^{(3)}$	
$\mathfrak{g}_{7,3.21}$	$\begin{pmatrix} 2 & -2 & -1 \\ -1 & 2 & 0 \\ -2 & 0 & 2 \end{pmatrix}$		$A_4^{(2)}$		
$\mathfrak{g}_{7,3.22}$	$\begin{pmatrix} 2 & -1 & -2 \\ -1 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}$				
$\mathfrak{g}_{7,3.23}$	$\begin{pmatrix} 2 & -2 & -1 \\ -2 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}$				
$\mathfrak{g}_{7,3.24}$	$\begin{pmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & -1 \\ 0 & -1 & 2 & -1 \\ 0 & -1 & -1 & 2 \end{pmatrix}$	A_4		$H_3^{(4)}$	
$\mathfrak{g}_{7,4.1}$	$\begin{pmatrix} 2 & -1 & -1 & 0 \\ -1 & 2 & 0 & 0 \\ -1 & 0 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{pmatrix}$				
$\mathfrak{g}_{7,4.2}$	$\begin{pmatrix} 2 & -1 & -1 & -1 \\ -1 & 2 & 0 & 0 \\ -1 & 0 & 2 & 0 \\ -1 & 0 & 0 & 2 \end{pmatrix}$	D_4			
$\mathfrak{g}_{7,4.3}$	$\begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & -1 \\ 0 & 0 & 0 & 2 & -1 \\ 0 & 0 & -1 & -1 & 2 \end{pmatrix}$	$A_4 \times A_3$			
$\mathfrak{g}_{7,4.4}$	$\begin{pmatrix} 2 & 0 & 0 & -1 & 0 & 0 \\ 0 & 2 & 0 & 0 & -1 & 0 \\ 0 & 0 & 2 & 0 & 0 & -1 \\ -1 & 0 & 0 & 2 & 0 & 0 \\ 0 & -1 & 0 & 0 & 2 & 0 \\ 0 & 0 & -1 & 0 & 0 & 2 \end{pmatrix}$	$A_2 \times A_2 \times A_2$			